

# Providing Palliative Care

## Oncology nurses' perceptions of their self-reported abilities

Catherine M. Mann, EdD, RN, CNS, CNE, Fatimah Rashed Aldossary, MS, RN, BS, and Suzanne S. Sullivan, PhD, MBA, RN, CHPN



**BACKGROUND:** Oncology nurses are pivotal in the provision of primary palliative care. However, nurses working in oncology settings, where the goal is often curative, may lack knowledge about and comfort in providing palliative care.

**OBJECTIVES:** The purpose of this study is to determine how oncology nurses perceive their ability to provide palliative care services to patients and their families.

**METHODS:** Using Benner's theory of skill acquisition as a basis for self-evaluation, this cross-sectional, descriptive pilot study surveyed oncology nurses.

**FINDINGS:** All participants reported ranges from capable to very capable in ability to provide precepts of palliative care. Results showed a small but statistically significant association between years of experience and perceived ability to honor patient and family preferences, provide patient and family support, appreciate demands placed on families, and provide supportive services to caregivers.

### KEYWORDS

palliative care; oncology nurses; caregivers; quality of life; cancer diagnosis

### DIGITAL OBJECT IDENTIFIER

10.1188/19.CJON.647-654

**RECEIVING A CANCER DIAGNOSIS IS A LIFE-ALTERING EXPERIENCE**, broadly affecting individuals and their family members as they adapt to and cope with physical symptoms, such as pain, and the emotional burdens of uncertainty and fear. The Oncology Nursing Society (2019) and the National Comprehensive Cancer Network (2019) recommend that an interprofessional team deliver comprehensive, patient- and family-centered palliative care starting at diagnosis, concurrent with treatment, and continuing throughout the cancer journey, including survivorship. Therefore, oncology nurses, as members of the interprofessional team, are pivotal to providing primary palliative care, including direct care, care coordination, and patient and family advocacy and education (National Coalition for Hospice and Palliative Care, 2018). Palliative care, an approach for caring for patients who are facing life-threatening illness and their families, aims to improve quality of life “through the prevention and relief of suffering by means of early identification and impeccable assessment and treatment of pain and other problems, physical, psychosocial and spiritual” (World Health Organization, 2019, para. 1). Palliative care is a holistic approach to care that may benefit all people affected by a cancer diagnosis, including patients' families and caregivers (National Cancer Institute, 2017).

Despite the ongoing support from national bodies and professional organizations, and the widespread availability of guidelines and resources, the adoption of palliative care remains limited. To support and expand palliative care, the National Academies of Sciences, Engineering, and Medicine (formerly known as the Institute of Medicine [IOM], 2015) supports a national strategy to improve access to palliative care and improve healthcare workforce training. That training includes building skills in pain and symptom management and care coordination (IOM, 2015; Meier et al., 2017).

### Background

As a complement to clinical oncology care, nurses who provide palliative care are confident in their abilities and can advocate for patients. Few studies have identified nurses' ability to provide palliative care. Therefore, there is a lack of studies about palliative care education in clinical oncology settings. An integrative review by Achora and Labrague (2019) about nurses' knowledge about and attitudes toward providing palliative care in a variety of settings suggested that nurses have positive attitudes toward palliative care, despite

a general lack of knowledge—specifically, knowledge related to pain and symptom management, psychological care, and philosophies of palliative care delivery.

The purpose of this pilot study is to determine how oncology nurses perceive their ability to provide elements of primary palliative care to patients and their families. In addition, related to palliative care competencies, this study explores the relationship between participants' self-reported perception of ability to provide precepts of palliative care, years of experience, and highest level of education.

**Methods**

This was a cross-sectional, descriptive pilot study of oncology nurses practicing at the Roswell Park Comprehensive Cancer Center in Buffalo, New York. The institute provides comprehensive cancer care to the region and has a palliative care team available for consultation.

**Sample**

A convenience sample of oncology nurses was obtained from a recruitment letter sent via the cancer institute's email system, which invited potential participants to follow a link to an online

**“Nurses who provide palliative care are confident in their abilities and can advocate for patients.”**

survey on SurveyMonkey®. Data were collected during a four-week period after obtaining approval from the institutional review board at the cancer center and Daemen College in Amherst, New York. Seventy-nine nurses self-selected to voluntarily participate in the study.

**Measurement**

To document the study sample's demographics, study participants provided their highest level of education and years of

**TABLE 1.** PRECEPTS OF PALLIATIVE CARE AND THEIR DECLARATIVE STATEMENTS AND RELATED DOMAINS

PRECEPT	STATEMENTS	DOMAINS
Precept 1: Respecting patient goals, preferences, and choices	<ul style="list-style-type: none"> <li>Statement 1: Identify and honor patient and family preference through careful attention to values, goals, and priorities, as well as cultural and spiritual perspectives.</li> <li>Statement 2: Assist patients in establishing goals of care by facilitating understanding of diagnosis and prognosis, clarifying priorities, promoting informed choices, and providing opportunities for negotiating care with providers.</li> </ul>	<ul style="list-style-type: none"> <li>Domain 1: Structure and process of care</li> <li>Domain 7: Care of the patient nearing end of life</li> <li>Domain 8: Ethical and legal aspects of care</li> </ul>
Precept 2: Providing comprehensive care	<ul style="list-style-type: none"> <li>Statement 3: Provide physical, psychological, social, and spiritual support to help the patient and family adapt to anticipated decline associated with advanced, progressive, incurable disease.</li> <li>Statement 4: Alleviate isolation through a commitment to nonabandonment, ongoing communication, and sustainment of relationships.</li> </ul>	<ul style="list-style-type: none"> <li>Domain 1: Structure and processes of care</li> <li>Domains 2, 3, 4, 5, and 6: Physical, social, spiritual, religious, and existential; and cultural aspects of care</li> <li>Domain 7: Care of the patient nearing end of life</li> </ul>
Precept 3: Utilizing the strengths of interdisciplinary resources	<ul style="list-style-type: none"> <li>Statement 5: Appreciate substantial physical, emotional, and economic demands placed on families caring for someone at home.</li> <li>Statement 6: Have knowledge, skills, experience, and provision of opportunity for ongoing education, professional support, and development.</li> </ul>	<ul style="list-style-type: none"> <li>Domain 1: Structure and processes of care</li> <li>Domain 7: Care of the patient nearing end of life</li> <li>Domain 8: Ethical and legal aspects of care</li> </ul>
Precept 4: Acknowledging and addressing caregiver concerns	<ul style="list-style-type: none"> <li>Statement 7: Appreciate substantial physical, emotional, and economic demands placed on families caring for someone at home.</li> <li>Statement 8: Provide concrete supportive services to caregivers, such as respite, around-the-clock availability of expert advice and support, grief counseling, personal care assistance, and referral to community resources.</li> </ul>	<ul style="list-style-type: none"> <li>Domain 1: Structure and processes of care</li> <li>Domain 7: Care of the patient nearing end of life</li> <li>Domain 8: Ethical and legal aspects of care</li> </ul>

**Note.** Based on information from National Coalition for Hospice and Palliative Care, 2018; Task Force on Palliative Care et al., 1998.

Downloaded on 07-22-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

experience. The study survey was based on four of the five precepts of palliative care. The precepts guide nursing care and are foundational to clinical practice guidelines. The precepts use eight declarative statements (S1–S8) (National Coalition for Hospice and Palliative Care, 2018; Task Force on Palliative Care, Last Acts Campaign, & Robert Wood Johnson Foundation, 1998) (see Table 1). Precepts 1–4 focus on nurses' perceived abilities at the bedside. Precept 5 was not included in the survey because it is related to organizational infrastructure.

Using Benner's (2001) novice to expert categories, for each of the eight statements, participants ranked their perceived ability to provide the precepts (on a Likert-type scale ranging from 1 [novice] to 5 [expert]). The five stages of skill development in nursing are as follows (Benner, 2001):

- Novice: those without any experience
- Advanced beginner: able to guide actions based on rules and experiences
- Competent: some feelings of mastery based on experience, which serve as guidelines to plan care
- Proficient: ability to view situations as a whole because they have multiple perspectives on the situation
- Expert: inherent intuitive grasp formed through a deep understanding of the situation

When adapting Benner's (2001) theory of skill acquisition, both novice and expert remained ranked as defined by Benner. A participant who indicated feeling somewhat capable was considered equivalent to an advanced beginner, capable was considered to be competent, and very capable was deemed proficient.

### Analysis

Descriptive statistics were used to evaluate data collected using IBM SPSS Statistics, version 25.0. Analysis of variance (ANOVA) determined whether there was a statistically significant difference between highest level of education and years of experience, and highest level of education and responses to the eight statements on the precepts (S1–S8). Likert-type responses, from 1 (novice) to 5 (expert), were treated as a continuous variable. In addition, the Student–Newman–Keuls stepwise multiple comparisons procedure was used as a secondary test to determine statistical significance (i.e., risk of a false-positive result) ( $p < 0.05$ ). Correlations, which measure the strength and direction of the relationship, ranging from +1 to –1, between highest level of education and each precept (S1–S8) are reported using Spearman's rho. Correlations between years of nursing experience and precepts (S1–S8) are reported using Spearman's rho and Pearson's  $r$ . Pearson's  $r$  is used to measure the relationship between two continuous variables, and Spearman's rho evaluates continuous or ordinal-level variables. The strength of correlation for both approaches are considered to be small (0.1 to 0.2), medium (0.3 to 0.4), and large (0.5 to 1) (Polit & Beck, 2014).

**TABLE 2.**  
SAMPLE CHARACTERISTICS (N = 79)

CHARACTERISTIC	n
<b>Highest level of education</b>	
Associate degree	25
Bachelor's degree	36
Master's degree	13
Other	5
<b>Years of experience</b>	
0–1	4
2–3	4
4–5	2
6–10	17
More than 10	51
Missing data	1

### Findings

Seventy-nine oncology nurses participated in the study. The majority of the participants ( $n = 51$ ) had more than 10 years of experience, and 10 participants had fewer than 5 years of experience; one participant chose not to report experience (see Table 2). Based on 2017 benchmarks, the National Council of State Boards of Nursing (2019) established that 42% of nurses had a bachelor's degree as qualification for first license. Of the study participants, 36 (46%) had earned a bachelor's degree and 13 (16%) had earned a master's degree.

#### Perceptions of Ability to Provide Precepts of Palliative Care

The majority of the participants reported their ability to provide the precepts in the range of capable to very capable for all eight statements (see Table 3). Honoring preferences (S1) was the highest-ranked item, with 57 respondents indicating that they perceived themselves to be very capable and 7 considering themselves to be expert ( $\bar{X} = 3.9$ ,  $SD = 0.59$ ). This statement (S1) supports participants' perceived ability to identify and honor preferences of patients and families through careful attention to their values, goals, and priorities, as well as their cultural and spiritual perspectives. In addition, demands placed on families (S7, precept 4) indicated that 44 oncology nurses perceived themselves as being very capable and 17 as being experts in their ability to appreciate the substantial physical, emotional, and economic demands placed on families ( $\bar{X} = 3.9$ ,  $SD = 0.75$ ). This statement

**TABLE 3.** PARTICIPANTS' PERCEPTIONS OF ABILITY TO PROVIDE PRECEPTS OF PALLIATIVE CARE BY BENNER'S NOVICE TO EXPERT CATEGORIES (N = 79)

PRECEPT AND STATEMENTS	NOVICE (1)	SOMEWHAT CAPABLE (2)	CAPABLE (3)	VERY CAPABLE (4)	EXPERT (5)	$\bar{X}$	SD
<b>Precept 1</b>							
S1: Honoring preferences	–	2	13	57	7	3.9	0.59
S2: Goals of care <sup>a</sup>	–	4	21	46	7	3.7	0.71
<b>Precept 2</b>							
S3: Providing support	–	4	19	47	9	3.8	0.72
S4: Alleviating isolation	1	2	21	46	9	3.8	0.74
<b>Precept 3</b>							
S5: Team approach	1	4	21	47	6	3.7	0.75
S6: Knowledge and development opportunity <sup>a</sup>	–	5	17	47	9	3.8	0.74
<b>Precept 4</b>							
S7: Demands placed on families	–	3	15	44	17	3.9	0.75
S8: Caregiver supportive services	6	10	35	24	4	3.2	0.97
<sup>a</sup> N = 78 S—statement Note. Based on information from Benner, 2001.							

(S7) relates to the participants' ability to acknowledge and address caregiver concerns.

**Years of Nursing Experience by Highest Level of Education**

There was no statistical difference in the sample between highest level of education and years of experience, except for participants without an academic degree. Five respondents had, on average, 27.2 years of experience but did not indicate having a conferred associate, bachelor's, or master's degree (see Tables 4 and 5).

**Perception of Ability**

There was a small but statistically significant positive correlation between years of nursing experience and honoring preferences (S1) ( $r = 0.23$ ), providing support (S3) ( $r = 0.28$ ), demands placed on families (S7) ( $r = 0.14$ ,  $p < 0.05$ ), and caregiver supportive services (S8) ( $r = 0.3$ ,  $p < 0.01$ ). Years of nursing experience and knowledge and development opportunity (S6) were near statistical significance ( $r = 0.28$ ,  $p = 0.06$ ), which suggests that there may be an effect but not enough of one to reject the null hypothesis. Universally, caregiver supportive services (S8) was the weakest perceived ability reported. There was a medium positive correlation between years of experience and caregiver supportive

services (S8) ( $r = 0.31$ ,  $p < 0.01$ ) (see Table 6). However, there was no statistically significant correlation between highest level of education and S1–S8.

**Discussion**

This pilot study reports on the perceived ability of oncology nurses to deliver the precepts of palliative care at a large urban inpatient cancer institute in New York. Overall, oncology nurses reported that they had a strong ability to provide the precepts; this finding supports the current clinical practice guidelines to deliver palliative care as an interprofessional team (National Coalition for Hospice and Palliative Care, 2018). This study revealed that years of nursing experience correlates with precepts of palliative care in several areas; however, highest level of education was not associated with any of the precepts. These findings suggest that, although nurses may perceive themselves as being proficient, this is not necessarily a product of years of formal education, but rather from experience or specific training in palliative care. Benner (2001) stated that the number of years of nursing experience in the same work environment, and in the same or similar medical situations, can establish and build competence in nurses' clinical practice. As revealed in this pilot study's results, nurses

Downloaded on 07-22-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

**TABLE 4.**  
DESCRIPTIVE STATISTICS FOR HIGHEST LEVEL  
OF EDUCATION RELATED TO YEARS OF EXPERIENCE  
AND PRECEPTS 1–4 (S1–S8) (N = 79)

CHARACTERISTIC	n	$\bar{X}$	SD	95% CI
<b>Years of experience</b>				
Associate degree	25	17.3	8.32	[13.91, 20.77]
Bachelor's degree	36	15.8	10.27	[12.26, 19.42]
Master's degree	13	14.8	9.9	[8.8, 20.74]
Other*	5	27.2	13.26	[10.74, 43.66]
<b>S1: Honoring preferences</b>				
Associate degree	25	3.6	0.57	[3.41, 3.87]
Bachelor's degree	36	4	0.54	[3.82, 4.18]
Master's degree	13	3.9	0.64	[3.54, 4.31]
Other	5	4	0.71	[3.12, 4.88]
<b>S2: Goals of care<sup>a</sup></b>				
Associate degree	25	3.64	0.64	[3.38, 3.9]
Bachelor's degree	36	3.75	0.78	[3.49, 4.01]
Master's degree	13	3.77	0.6	[3.41, 4.13]
Other	4	3.75	0.96	[2.23, 5.27]
<b>S3: Providing support</b>				
Associate degree	25	3.56	0.58	[3.32, 3.8]
Bachelor's degree	36	3.97	0.7	[3.74, 4.21]
Master's degree	13	3.77	0.73	[3.33, 4.21]
Other	5	3.4	1.14	[1.98, 4.82]
<b>S4: Alleviating isolation</b>				
Associate degree	25	3.76	0.66	[3.49, 4.03]
Bachelor's degree	36	3.81	0.79	[3.54, 4.07]
Master's degree	13	3.69	0.75	[3.24, 4.15]
Other	5	3.6	0.89	[2.49, 4.71]
<b>S5: Team approach</b>				
Associate degree	25	3.56	0.65	[3.29, 3.83]
Bachelor's degree	36	3.72	0.78	[3.46, 3.99]
Master's degree	13	3.69	0.9	[3.18, 4.21]

*Continued in the next column*

**TABLE 4. (CONTINUED)**  
DESCRIPTIVE STATISTICS FOR HIGHEST LEVEL  
OF EDUCATION RELATED TO YEARS OF EXPERIENCE  
AND PRECEPTS 1–4 (S1–S8) (N = 79)

CHARACTERISTIC	n	$\bar{X}$	SD	95% CI
<b>S5: Team approach (continued)</b>				
Other	5	3.8	0.84	[2.76, 4.84]
<b>S6: Knowledge and development opportunity<sup>a</sup></b>				
Associate degree	24	3.58	0.65	[3.31, 3.86]
Bachelor's degree	36	3.83	0.74	[3.58, 4.08]
Master's degree	13	4	0.81	[3.51, 4.49]
Other	5	3.6	0.89	[2.49, 4.71]
<b>S7: Demands placed on families</b>				
Associate degree	25	3.72	0.71	[3.42, 4.02]
Bachelor's degree	36	4.03	0.74	[3.78, 4.28]
Master's degree	13	4.15	0.69	[3.74, 4.57]
Other	5	4	1	[2.76, 5.24]
<b>S8: Caregiver supportive services</b>				
Associate degree	25	3.12	0.73	[2.82, 3.42]
Bachelor's degree	36	3.17	1.03	[2.82, 3.51]
Master's degree	13	3.31	0.9	[2.79, 3.82]
Other	5	2.4	1.67	[0.32, 4.48]

\* p < 0.05

<sup>a</sup> N = 78

CI—confidence interval; S—statement

**Note.** Mean years of nursing experience was significantly different for nurses without formal degrees (licensed practical nurse, n = 1; diploma in nursing, n = 2; 5 years of college, n = 1; some graduate school, n = 1), as determined by the Student–Newman–Keuls procedure.

with additional years of experience perceived their abilities as higher than their less experienced counterparts. In addition, these findings may also be influenced by exposure to, and interactions with, the palliative care consultation team.

This pilot study's findings support the findings from previous studies about nurses' years of experience and their perceptions of their ability to provide palliative and end-of-life care. For example, Price et al. (2017) reported that nurses with more than 10 years of experience working in intensive care perceived themselves to be more competent in providing end-of-life care, scoring

higher knowledge, decision making, emotional support for family, and spiritual support competencies than nurses with fewer than 10 years of experience. In the current study, the majority of participants had more than 10 years of experience and reported themselves as being capable or very capable in their ability to perform precepts of palliative care, as measured in S1–S7.

The lowest mean score for perceived ability was in precept 4, caregiver supportive services (S8); however, this statement was most highly correlated with years of nursing experience, suggesting that more experienced nurses are more comfortable in providing support to patients and their families. In addition, nurses who practice in oncology settings are highly attuned to meeting the physical and emotional needs of patients with cancer, regardless of whether they are receiving palliative care services, in contrast to nurses in general practice who identify having poor knowledge of approaches for pain and symptom management (Achora & Labrague, 2019). Therefore, this may have an impact on the participants' report of expertise in precept 2, which is related to the physical and emotional demands of patients.

**Limitations**

Pilot study participants volunteered to participate in the study and provided self-reported perceptions. This could influence the results because of self-selection and variations in personal practice patterns. In addition, the small sample size limits generalizability to inpatient oncology settings. The evaluation tool was developed by the researchers, and it has not been validated. This pilot study did not address a complete set of clinical practice

**TABLE 5.**  
ANOVA SUMMARY OF YEARS OF EXPERIENCE AND S1–S8 (N = 79)

CHARACTERISTIC	F	df	p
Years of experience	2.07	3	0.11
S1: Honoring preferences	2.07	3	0.11
S2: Goals of care <sup>a</sup>	0.15	3	0.93
S3: Providing support	2.2	3	0.1
S4: Alleviating isolation	0.167	3	0.92
S5: Team approach	0.28	3	0.84
S6: Knowledge and development opportunity <sup>a</sup>	1.15	3	0.33
S7: Demands placed on families	1.25	3	0.3
S8: Caregiver supportive services	1.13	3	0.34

<sup>a</sup>N = 78  
ANOVA—analysis of variance; df—degrees of freedom; S—statement

**TABLE 6.**  
CORRELATIONS OF PRECEPTS WITH HIGHEST LEVEL OF EDUCATION AND YEARS OF EXPERIENCE

STATEMENT	HIGHEST LEVEL OF EDUCATION	YEARS OF EXPERIENCE	
	rho <sup>a</sup>	rho <sup>a</sup>	r <sup>b</sup>
S1: Honoring preferences	0.19	0.16	0.23**
S2: Goals of care	0.09	0.07	0.13
S3: Providing support	0.2	0.21	0.28**
S4: Alleviating isolation	-0.06	0.16	0.19
S5: Team approach	0.09	0.13	0.22
S6: Knowledge and development opportunity	0.19	0.21***	0.28
S7: Demands placed on families	0.19	0.13	0.14**
S8: Caregiver supportive services	0.06	0.31*	0.3*

\*p < 0.01; \*\* p < 0.05; \*\*\* p = 0.06  
<sup>a</sup>Spearman's rho rank correlation coefficient  
<sup>b</sup>Pearson's linear correlation coefficient  
**Note.** Strength of association is as follows: 0 (no association), 0.1 to 0.2 (small association), 0.3 to 0.4 (medium association), and 0.5 to 1 (large association).

guidelines because the study focused on selected precepts of palliative care.

In addition, the “other” group in years of experience combined a small sample of individuals with conceptually unrelated levels of experience, which could affect the findings of this pilot study. There is also a wide 95% confidence interval in this domain, which reduces the certainty of the finding. However, it was necessary to group these individuals together for analysis because there were only five participants without a traditional nursing education.

**Implications for Practice**

Based on this pilot study's findings, nurses with more than 10 years of experience, who are defined as experts in Benner's (2001) model, may act as guides and facilitators for novice and advanced beginners in nursing who are learning how to provide palliative care. Clinical oncology nurses have the expertise to manage patients with cancer, supporting them and their family members during the trajectory of what may be incurable disease. This expertise includes delivering palliative care. However, nurses may not have the confidence and experience related to caregiver supportive services (S8, precept 4). These findings emphasize the importance of education for nurses in practice and students in nursing programs so that they can improve their abilities and confidence in supporting patients' families. Initiatives that support palliative care nurse education can significantly affect



nurses' foundational knowledge about palliative care, as well as their attitudes toward and behaviors in better delivering that care (Harden, Price, Duffy, Galunas, & Rodgers, 2017).

This study's findings endorse the call from the American Association of Colleges of Nursing's new competencies and recommendations for educating undergraduate nursing students, which urge nurse educators to implement 17 palliative care competencies in undergraduate curricula (Ferrell, Malloy, Mazanec, & Virani, 2016). Several of these nursing competencies are family-centered, including educating family about palliative care issues (competency 5); collaborating with members of the interprofessional team, specifically aimed at ensuring coordinated care (competency 6); assisting the family to cope and build resilience (competency 16); and recognizing the need to seek consultation (competency 17) (Ferrell et al., 2016).

The current study also supports palliative care delivered by interprofessional teams, particularly to address caregiver needs. To advance interprofessional team care, experienced nurses can champion innovative methods to build competencies among team members beyond formal education programs. These methods may include focused mentorship and simulations.

## Conclusion

Based on this pilot study's findings, oncology nurses consistently have opportunities to develop their skills in primary palliative nursing; these skills are gained through time and experience caring for patients with serious illness and their families. Engaging oncology nurses, particularly those with more than 10 years of experience—in mentorship, preceptor, and other roles in nursing education—may help bolster comfort and competence in providing palliative care. As members of interprofessional palliative care teams, experienced nurses can provide care to caregivers, ensuring that they have access to supportive services.

**Catherine M. Mann, EdD, RN, CNS, CNE**, is the assistant dean of undergraduate studies and a clinical associate professor of nursing in the School of Nursing at the University of Buffalo in New York; **Fatimah Rashed Aldossary, MS, RN, BS**, is the postgraduate oncology nursing diploma program director at King Fahad Specialist Hospital-Dammam in Saudi Arabia; and **Suzanne S. Sullivan, PhD, MBA, RN, CHPN**, is an assistant professor in the School of Nursing at the University of Buffalo. Mann can be reached at [cmmann@buffalo.edu](mailto:cmmann@buffalo.edu), with copy to [CJONEditor@ons.org](mailto:CJONEditor@ons.org). (Submitted April 2019. Accepted June 13, 2019.)

The authors take full responsibility for this content and did not receive honoraria or disclose any relevant financial relationships. The article has been reviewed by independent peer reviewers to ensure that it is objective and free from bias.

## REFERENCES

- Achora, S., & Labrague, L.J. (2019). An integrative review on knowledge and attitudes of nurses toward palliative care: Implications for practice. *Journal of Hospice and Palliative Nursing*, *21*, 29–37. <https://doi.org/10.1097/NJH.0000000000000481>

## IMPLICATIONS FOR PRACTICE

- Improve abilities to provide primary palliative care by accessing professional development programs, such as the End-of-Life Nursing Education Consortium.
- Collaborate with and seek mentorship from experienced nurses to improve palliative care competencies.
- Promote patient- and family-centered care in oncology care settings.

- Benner, P.E. (2001). *From novice to expert: Excellence and power in clinical nursing practice* (Commemorative ed.). Upper Saddle River, NJ: Prentice Hall.
- Ferrell, B., Malloy, P., Mazanec, P., & Virani, R. (2016). CARES: AACN's new competencies and recommendations for educating undergraduate nursing students to improve palliative care. *Journal of Professional Nursing*, *32*, 327–333. <https://doi.org/10.1016/j.profnurs.2016.07.002>
- Harden, K., Price, D., Duffy, E., Galunas, L., & Rodgers, C. (2017). Palliative care: Improving nursing knowledge, attitudes, and behaviors [Online exclusive]. *Clinical Journal of Oncology Nursing*, *21*, E232–E238. <https://doi.org/10.188/17CJON.E232-E238>
- Institute of Medicine. (2015). *Dying in America: Improving quality and honoring individual preferences near the end of life*. Washington, DC: National Academies Press.
- Meier, D.E., Back, A.L., Berman, A., Block, S.D., Corrigan, J.M., & Morrison, R.S. (2017). A

## PROFESSIONAL DEVELOPMENT ACTIVITY EARN 0.5 CONTACT HOURS



ONS members can earn free nursing continuing professional development (NCPD) for reading this article and completing an evaluation online. To do so, visit [cjon.ons.org/ncpd](http://cjon.ons.org/ncpd) to link to this article and then access its evaluation link after logging in.

Certified nurses can claim no more than 0.5 total ILNA points for this program. Up to 0.5 ILNA points may be applied to Palliative Care OR Psychosocial OR Oncology Nursing Practice OR Professional Practice. See [www.oncc.org](http://www.oncc.org) for complete details on certification.

## QUESTIONS FOR DISCUSSION USE THIS ARTICLE FOR JOURNAL CLUB



Journal club programs can help to increase your ability to evaluate the literature and translate those research findings to clinical practice, education, administration, and research. Use the following questions to start the discussion at your next journal club meeting.

- What do you think about your ability to provide primary palliative care to patients? What do you do well? What skills do you want to improve?
- So that nurses new to the profession can learn from key experiences, what experiences do you recommend that best build primary palliative care competencies?
- What was a situation in which you taught that caregiver a specific palliative care skill? How did you know your teaching strategy was effective (or not) when teaching the caregiver?

Visit <http://bit.ly/1vUqBvj> for details on creating and participating in a journal club. Photocopying of this article for discussion purposes is permitted.

- national strategy for palliative care. *Health Affairs*, 36, 1265–1273. <https://doi.org/10.1377/hlthaff.2017.0164>
- National Cancer Institute. (2017). Palliative care in cancer. Retrieved from <https://www.cancer.gov/about-cancer/advanced-cancer/care-choices/palliative-care-fact-sheet>
- National Coalition for Hospice and Palliative Care. (2018). *Clinical practice guidelines for quality palliative care* (4th ed.). Richmond, VA: Author.
- National Comprehensive Cancer Network. (2019). *NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®): Palliative care* [v.2.2019]. Retrieved from [https://www.nccn.org/professionals/physician\\_gls/PDF/palliative.pdf](https://www.nccn.org/professionals/physician_gls/PDF/palliative.pdf)
- National Council of State Boards of Nursing. (2019). National nursing workforce study. Retrieved from <https://www.ncsbn.org/workforce.htm>
- Oncology Nursing Society. (2019). Palliative care for people with cancer. Retrieved from <https://www.ons.org/make-difference/ons-center-advocacy-and-health-policy/position-statements/palliative-care-people>
- Polit, D.F., & Beck, C.T. (2014). *Essentials of nursing research: Appraising evidence for nursing practice* (8th ed.). Philadelphia, PA: Wolters Kluwer.
- Price, D.M., Strodman, L., Montagnini, M., Smith, H.M., Miller, J., Zybert, J., . . . Ghosh, B. (2017). Palliative and end-of-life care education needs of nurses across inpatient care settings. *Journal of Continuing Education in Nursing*, 48, 329–336.
- Task Force on Palliative Care, Last Acts Campaign, & Robert Wood Johnson Foundation. (1998). Precepts of palliative care. *Journal of Palliative Medicine*, 1, 109–112. <https://doi.org/10.1089/jpm.1998.1.109>
- World Health Organization. (2019). WHO definition of palliative care. Retrieved from <https://www.who.int/cancer/palliative/definition/en>