

Appendix. Cancer treatment-related lymphedema

Jingyi Frances Ding, MD

Bashar Hasan, MD

Konstantinos Malandris, MD, MSc

Magdoleen H. Farah, MBBS

Apostolos Manolopoulos, MD, MSc

Pamela Ginex, EdD, RN

Allison B. Anbari, PhD, RN

Tarek Nayfeh, MD

Moutie R. Rajjoub

Raed Benkhadra, MD

Larry J. Prokop, MLS

Rebecca L. Morgan, PhD, MPH

M. Hassan Murad, MD, MPH

Table of Contents

Search strategy	3
Figure S1. Risk of lymphedema (other interventions).....	9
Figure S2. Change in physical activity	10
Figure S3. Grip strength.....	11
Figure S4. Range of motion	12
Figure S5. Risk of bias assessment for randomized clinical trials	16
Figure S6. Risk of bias assessment for observational studies and non-randomized clinical trials	17
Table S1. Protocol of cancer related lymphedema guideline and systematic review.....	18
Table S2. Characteristics of studies in the meta-analysis by comparison	19
Table S3. Certainty of evidence of the studies included in the meta-analysis.....	38
Table S4. Characteristics of included studies comparing Progressive resistance training (RRT) vs Usual care	43
Table S5. Characteristics of included studies comparing Compression vs No compression	45
Table S6. Characteristics of included studies comparing Manual lymph drainage (MLD) vs No MLD	46
Table S7. Characteristics of included studies comparing Manual lymph drainage (MLD) vs Exercise.....	51
Table S8. Characteristics of included studies comparing Exercise vs No exercise.....	53
Table S9. Characteristics of included studies comparing Delayed exercise vs Early exercise.....	61
Table S10. Characteristics of included studies comparing Physiotherapy vs No physiotherapy	63
Table S11. Characteristics of included studies comparing Surveillance, education, or clinical care vs No intervention or standard care.....	66
References.....	70

Search strategy

Ovid

Database(s): EBM Reviews - Cochrane Central Register of Controlled Trials February 2019, EBM Reviews - Cochrane Database of Systematic Reviews 2005 to March 27, 2019, Embase 1974 to 2019 March 29, Ovid MEDLINE(R) and Epub Ahead of Print, In-Process & Other Non-Indexed Citations and Daily 1946 to March 29, 2019

Search Strategy:

#	Searches	Results
1	exp Neoplasms/rt, su [Radiotherapy, Surgery] (cancer* or neoplasm* or neoplastic or paraneoplas* or tumor* or tumor* or neoplasia* or "section 16" or leukemia* or carcinoma* or lymphoma* or Astrocytoma* or glioma* or adenoma* or carcinoid* or Sarcoma* or ostesarcoma* or histiocytoma* or craniopharyngioma* or ependymoma* or chordoma* or "Chronic Myeloproliferative Disorder*" or craniopharyngioma* or "Mycosis Fungoide*" or "Sézary Syndrome*" or Esthesioneuroblastoma* or melanoma* or retinoblastoma* or histeocytoma* or "gestational trophoblastic disease*" or histiocytos* or burkitt* or Macroglobulinemia* or Mesothelioma* or neuroblastoma* or Papillomatos* or paraganglioma* or "pheochromocytoma* multiple myeloma*" or blastoma* or Rhabdomyosarcoma* or nonmelanoma* or metasta*).ti,ab,hw,kw.	1391485
2		9341523
3	exp Radiotherapy/	648747
4	exp surgical procedures, operative/	7468924
5	(Brachytherap* or irradiat* or operat* or radiation* or "radio therap*" or "radio treatment*" or radiolog* or radiotherap* or radiotreatment* or reconstruction* or repair* or resect* or roentgen* or rontgen* or surg* or "x ray*").ti,ab,hw,kw.	10191551
6	3 or 4 or 5	13342834
7	1 or (2 and 6)	3626644
8	exp Lymphedema/ ("bigfoot disease*" or elephantiasis or elephantiasis or "lymph edema*" or "lymph oedema*" or "lymphatic edema*" or "lymphatic filariases" or "lymphatic filariasis" or "lymphatic oedema*" or lymphedema* or lymphoedema* or lymphooedema* or "lymphostatic edema*" or "lymphostatic oedema*" or "lymphostatic verrucoses" or "lymphostatic verrucosis" or "microcrystal disease*" or "milroy disease" or "milroy meige disease" or "milroy meiges disease" or "milroys disease" or "mossy feet" or "mossy foot*" or podoconioses or podoconiosis).ti,ab,hw,kw.	30546
9		38766
10	8 or 9	39158
11	7 and 10	11153
12	exp DISEASE SURVEILLANCE/	24791
13	(exam or examination* or exams or "patient report outcome*" or "patient reported outcome*" or PROM or surveillance).ti,ab,hw,kw.	2779044
14	exp Massage/	19943

15 ("manual lymph drainage" or "manual lymphatic drainage" or massag*).ti,ab,hw,kw.	38848
16 exp exercise/	505444
17 exp exercise therapy/	127546
18 (aerobics or anaerobics or bicycling or biking or "endurance training" or exercis* or "fitness training" or isometrics or "physical exertion" or "physical activit*" or "resistance training" or running or "strength training" or swimming or walking or weightlifting).ti,ab,hw,kw.	1432369
19 exp Stockings, Compression/	4050
20 exp compression therapy/	8299
21 ("circ aid" or circaid or compression or unna or unnas).ti,ab,hw,kw.	284949
22 or/12-21	4419563
23 11 and 22	2910
24 exp evidence based medicine/	1145422
25 exp meta analysis/	258169
26 exp Meta-Analysis as Topic/	57039
27 exp "systematic review"/	301029
28 exp Guideline/ or exp Practice Guideline/	522025
29 exp controlled study/	6680981
30 exp Randomized Controlled Trial/	1020185
31 exp triple blind procedure/	204
32 exp Double-Blind Method/	439387
33 exp Single-Blind Method/	79905
34 exp latin square design/	361
35 exp Placebos/	388945
36 exp Placebo Effect/	11206
37 exp comparative study/	3128292
38 exp intervention studies/	39732
39 exp Cross-Sectional Studies/	586142
40 exp Cross-Over Studies/	138371
41 exp Cohort Studies/	2429563

42	exp longitudinal study/	377741
43	exp retrospective study/	1496634
44	exp prospective study/	1090455
45	exp clinical trial/	2198408
46	clinical study/	156767
47	exp case-control studies/	1149203
48	exp confidence interval/	164691
49	exp multivariate analysis/ ((evidence adj based) or (meta adj analys*) or (systematic* adj3 review*) or guideline* or (control* adj3 study) or (control* adj3 trial) or (randomized adj3 study) or (randomized adj3 trial) or (randomised adj3 study) or (randomised adj3 trial) or "pragmatic clinical trial" or (doubl* adj blind*) or (doubl* adj mask*) or (singl* adj blind*) or (singl* adj mask*) or (tripl* adj blind*) or (tripl* adj mask*) or (trebl* adj blind*) or (trebl* adj mask*) or "latin square" or placebo* or nocebo* or multivariate or "comparative study" or "comparative survey" or "comparative analysis" or (intervention* adj2 study) or (intervention* adj2 trial) or "cross-sectional study" or "cross-sectional analysis" or "cross-sectional survey" or "cross-sectional design" or "prevalence study" or "prevalence analysis" or "prevalence survey" or "disease frequency study" or "disease frequency analysis" or "disease frequency survey" or crossover or "cross-over" or cohort* or "longitudinal study" or "longitudinal survey" or "longitudinal analysis" or "longitudinal evaluation" or longitudinal* or ((retrospective or "ex post facto") adj3 (study or survey or analysis or design)) or retrospectiv* or "prospective study" or "prospective survey" or "prospective analysis" or prospectiv* or "concurrent study" or "concurrent survey" or "concurrent analysis" or "clinical study" or "clinical trial" or "case control study" or "case base study" or "case referrent study" or "case referent study" or "case referent study" or "case compeer study" or "case comparison study" or "matched case control" or "multicenter study" or "multi-center study" or "odds ratio" or "confidence interval" or "change analysis" or ((study or trial or random* or control*) and compar*)).mp.pt.	514350
50		21782897
51	or/24-50	22337008
52	23 and 51 limit 52 to ("all adult (19 plus years)" or "young adult (19 to 24 years)" or "adult (19 to 44 years)" or "young adult and adult (19-24 and 19-44)" or "middle age (45 to 64 years)" or "middle aged (45 plus years)" or "all aged (65 and over)" or "aged (80 and over)") [Limit not valid in CCTR,CDSR,Embase; records were retained]	1682
54	limit 53 to (adult <18 to 64 years> or aged <65+ years>) [Limit not valid in CCTR,CDSR,Ovid MEDLINE(R),Ovid MEDLINE(R) Daily Update,Ovid MEDLINE(R) In-Process,Ovid MEDLINE(R) Publisher; records were retained]	1043
55	limit 52 to ("all infant (birth to 23 months)" or "all child (0 to 18 years)" or "newborn infant (birth to 1 month)" or "infant (1 to 23 months)" or "preschool child (2 to 5 years)" or "child (6 to 12 years)" or "adolescent (13 to 18 years)") [Limit not valid in CCTR,CDSR,Embase; records were retained]	1239
56	limit 55 to (embryo or infant or child or preschool child <1 to 6 years> or school child <7 to 12 years> or adolescent <13 to 17 years>) [Limit not valid in 227	

CCTR,CDSR,Ovid MEDLINE(R),Ovid MEDLINE(R) Daily Update,Ovid MEDLINE(R) In-Process,Ovid MEDLINE(R) Publisher; records were retained]

57 56 not 54	12
58 52 not 57	1670
59 ("case study" or "case series" or "clinical series" or "case studies" or (case adj3 report)).mp,pt.	3146591
60 58 not 59	1603
61 limit 60 to (editorial or erratum or note or addresses or autobiography or bibliography or biography or blogs or comment or dictionary or directory or interactive tutorial or interview or lectures or legal cases or legislation or news or newspaper article or overall or patient education handout or periodical index or portraits or published erratum or video-audio media or webcasts) [Limit not valid in CCTR,CDSR,Embase,Ovid MEDLINE(R),Ovid MEDLINE(R) Daily Update,Ovid MEDLINE(R) In-Process,Ovid MEDLINE(R) Publisher; records were retained]	20
62 from 61 keep 1-3	3
63 (60 not 61) or 62	1586
64 remove duplicates from 63	1130

Scopus

- 1 TITLE-ABS-KEY(cancer* or neoplasm* or neoplastic or paraneoplas* or tumor* or tumor* or neoplasia* or "section 16" or leukemia* or carcinoma* or lymphoma* or Astrocytoma* or glioma* or adenoma* or carcinoid* or Sarcoma* or osteosarcoma* or histiocytoma* or craniopharyngioma* or ependymoma* or chordoma* or "Chronic Myeloproliferative Disorder*" or craniopharyngioma* or "Mycosis Fungoide*" or "Szary Syndrome*" or Esthesioneuroblastoma* or melanoma* or retinoblastoma* or histiocytoma* or "gestational trophoblastic disease*" or histiocytos* or burkitt* or Macroglobulinemia* or Mesothelioma* or neuroblastoma* or Papillomatos* or paraganglioma* or "pheochromocytoma* multiple myeloma*" or blastoma* or Rhabdomyosarcoma* or nonmelanoma* or metasta*)
- 2 TITLE-ABS-KEY(Brachytherap* OR irradiat* OR operat* OR radiation* OR "radio therap*" OR "radio treatment*" OR radiolog* OR radiotherap* OR radiotreatment* OR reconstruction* OR repair* OR resect* OR roentgen* OR rontgen* OR surg* OR "x ray*")
- 3 TITLE-ABS-KEY("bigfoot disease*" OR elephantiasis OR elephantiasis OR "lymph edema*" OR "lymph oedema*" OR "lymphatic edema*" OR "lymphatic filariases" OR "lymphatic filariasis" OR "lymphatic oedema*" OR lymphedema* OR lymphoedema* OR lymphoedema* OR "lymphostatic edema*" OR "lymphostatic oedema*" OR "lymphostatic verrucoses" OR "lymphostatic verrucosis" OR "microcrystal disease*" OR "milroy disease" OR "milroy meige disease" OR "milroy meiges disease" OR "milroys disease" OR "mossy feet" OR "mossy foot*" OR podoconioses OR podoconiosis)
- 4 TITLE-ABS-KEY(exam or examination* or exams or "patient report outcome*" or "patient reported outcome*" or PROM or surveillance)
- 5 TITLE-ABS-KEY("manual lymph drainage" OR "manual lymphatic drainage" OR massag*)
- 6 TITLE-ABS-KEY(aerobics or anaerobics or bicycling or biking or "endurance training" or exercis* or "fitness training" or isometrics or "physical exertion" or "physical activit*" or "resistance training" or running or "strength training" or swimming or walking or weightlifting)
- 7 TITLE-ABS-KEY("circ aid" OR circaid OR compression OR unna OR unnas)
- 8 TITLE-ABS-KEY((evidence W/1 based) or (meta W/1 analys*) or (systematic* W/3 review*) or guideline* or (control* W/3 study) or (control* W/3 trial) or (randomized W/3 study) or (randomized W/3 trial) or (randomised W/3 study) or (randomised W/3 trial) or "pragmatic clinical trial" or (doubl* W/1 blind*) or (doubl* W/1 mask*) or (singl* W/1 blind*) or (singl* W/1 mask*) or (tripl* W/1 blind*) or (tripl* W/1 mask*) or (trebl* W/1 blind*) or (trebl* W/1 mask*) or "latin square" or placebo* or nocebo* or multivariate or "comparative study" or "comparative survey" or "comparative analysis" or (intervention* W/2 study) or (intervention* W/2 trial) or "cross-sectional study" or "cross-sectional analysis" or "cross-sectional survey" or "cross-sectional design" or "prevalence study" or "prevalence analysis" or "prevalence survey" or "disease frequency study" or "disease frequency analysis" or "disease frequency survey" or crossover or "cross-over" or cohort* or "longitudinal study" or "longitudinal survey" or "longitudinal analysis" or "longitudinal evaluation" or longitudinal* or ((retrospective or "ex post facto") W/3 (study or survey or analysis or design)) or retrospectiv* or "prospective study" or "prospective survey" or "prospective analysis" or prospectiv* or "concurrent study" or "concurrent survey" or "concurrent analysis" or "clinical study" or "clinical trial" or "case control study" or "case base study" or "case referent study" or "case referent study" or "case referent study" or "case compeer study" or "case comparison study" or "matched case control" or "multicenter study" or "multi-center study" or "odds ratio" or "confidence interval" or "change analysis" or ((study or trial or random* or control*) and compar*))
- 9 1 and 2 and 3 and (4 or 5 or 6 or 7) and 8
- 10 TITLE-ABS-KEY(newborn* or neonat* or infant* or toddler* or child* or adolescent* or paediatric* or pediatric* or girl or girls or boy or boys or teen or teens or teenager* or preschooler* or "pre-schooler*" or preteen or preteens or "pre-teen" or "pre-teens" or youth or youths) AND NOT TITLE-ABS-KEY(adult or adults or "middle age" or "middle aged" OR elderly OR geriatric* OR "old people" OR "old person*" OR "older people" OR "older person*" OR "very old")

- 11 9 and not 10
- 12 TITLE-ABS-KEY("case study" OR "case series" OR "clinical series" OR "case studies" OR (case W/3 report))
- 13 11 and not 12
- 14 DOCTYPE(ed) OR DOCTYPE(bk) OR DOCTYPE(er) OR DOCTYPE(no) OR DOCTYPE(sh)
- 15 13 and not 14
- 16 INDEX(embase) OR INDEX(medline) OR PMID(0* OR 1* OR 2* OR 3* OR 4* OR 5* OR 6* OR 7* OR 8* OR 9*)
- 17 15 and not 16

Figure S1. Risk of lymphedema (other interventions)




Figure S2. Change in physical activity

(Data are standardized. Higher SMD means improvement in the intervention group)




Figure S3. Grip strength






Figure S4. Range of motion


A. Shoulder flexion



B. Shoulder abduction



C. Shoulder external rotation



D. Shoulder horizontal extension




Figure S5. Risk of bias assessment for randomized clinical trials




Figure S6. Risk of bias assessment for observational studies and non-randomized clinical trials




Table S1. Protocol of cancer related lymphedema guideline and systematic review

Population of interest	Intervention	Comparison	Outcomes	Study designs available to answer this question	Timing
Adult patients who have had cancer surgery and/or radiation and are at risk for lymphedema	<ol style="list-style-type: none"> 1. Prospective surveillance, timing 2. Prospective surveillance components/assessment (objective measurements, PROM and clinical examination) 	No surveillance	<ol style="list-style-type: none"> 1. Development of lymphedema 2. Missed work or cost of coming in for prospective surveillance 3. # of patients referred to lymphedema specialist (surrogate for development of lymphedema) 	<p>Comparative observational studies</p> <p>RCT</p>	Months
Adult patients with cancer who are at risk for extremity/trunk lymphedema	<ol style="list-style-type: none"> 1. Prophylactic massage of scar tissue (by patient) 2. Preventive exercises 3. Prophylactic use of compression garments 	No treatment or an alternative intervention	<ol style="list-style-type: none"> 1. Development of persistent vs transient stage of lymphedema 2. Change in physical activity 3. Functional impairments (ROM, grip) 	<p>Comparative observational studies</p> <p>RCT</p>	Months

Table S2. Characteristics of studies in the meta-analysis by comparison

A. Compression vs. No compression

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Stuiver, 2013 (Stuiver et al., 2013)	RCT	80 (41/39)	59/58	49/59	NR	Melanoma/Ur ogenital Cancer Intervention: 22/18, control: 23/17	Surgery with inguinal lymph node dissection	NR	Compression stockings	Patient education	12	Risk of lymphedema: lower limb/genital, defined as a 10% or greater increase from baseline in volume of the proximal or distal half of the thigh or the lower leg, using the standardized Kühnke's method of surface measurement

B. MLD vs. No MLD

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Cho, 2016 (Cho et al., 2016)	RCT	41 (21/20)	46.6/50.7	0	NR	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Chemotherapy- Intervention:9 (42.9), control:11 (55). Radiotherapy- intervention:21 (100), control:19 (95).	Stage I: intervention:5 (23.8), control:12 (60) III: intervention:16 (76.2), control:8 (40).	Manual lymph drainage (MLD)+physical therapy (PT)	Physical therapy (PT)	1	1. Risk of lymphedema: ≥ 3 % volume increase from baseline in the affected upper limb 2. Volume change: arm volume calculated from circumference measurements taken at 4-cm intervals from the dorsum of the wrist to the axilla 3. Change in physical activity: EORTC QLQ-C30 and DASH 4. Functional impairment-range of motion: shoulder flexion/ abduction

Devoogdt, 2011 (Devoogdt et al., 2011)	RCT	160 (79/81)	55.8/54.5	1/1	26.6/26.2	Breast cancer	<p>Surgery- Mastectomy: Intervention: 52, Control: 56; Breast conserving: Intervention: 27, Control: 25. Radiotherapy- IMC and medial supraclavicular: Intervention: 69, Control: 67. Radiotherapy- axilla: Intervention: 8, Control: 5. Chemotherapy: Intervention: 50, Control: 58. Neo-adjuvant chemotherapy: Intervention: 14, Control: 14. Trastuzumab: Intervention: 14, Control: 7. Endocrine treatment: Intervention: 55, Control: 66.</p>	<p>Tumor size: pT0: intervention: 1, control: 0; pT1: intervention: 21, control: 26; pT2: intervention: 38, control:39; pT3: intervention: 13, control: 12; pT4: intervention: 6, control: 4.</p>	Manual lymph drainage (MLD)	No MLD	12	<p>1. Risk of lymphedema : (1) ≥ 200 mL increase in arm volume up to the reference point 16 cm proximal to the olecranon by water displacement (2) ≥ 2 cm increase in arm circumference at the olecranon and 4, 8, 12, 16, and 20 cm above and under the olecranon (3) using author developed questionnaire 2. Volume change : arm volume change 3. Change in physical activity : SF-36 (QoL : physical health)</p>
---	-----	-------------	-----------	-----	-----------	---------------	--	---	-----------------------------	--------	----	---

Devoogdt, 2018 (Devoogdt et al., 2018)	RCT (the same trial as Devoogdt, 2011)	160 (79/81)	55.8/54.5	1/1	26.6/26.2	Breast cancer	<p>All patients received surgery and some patients also received chemotherapy or radiation.</p> <p>Radiotherapy- intramammary chain, medial supraclavicular: intervention:69 (87), control:67 (83); axilla: intervention:8 (10), control:5 (6).</p> <p>Chemotherapy- all: intervention:50 (63), control:58 (72); neo-adjuvant: intervention:14 (18), control:14 (17). taxane-based: intervention:45 (57), control:46 (57).</p>	<p>Only for tumor size and node stage.</p> <p>Tumor size, n (%)</p> <p>pT0: intervention:1 (1) control:0 (0)</p> <p>pT1: intervention:21 (27) control:26 (32)</p> <p>pT2: intervention:38 (48) control:39 (48)</p> <p>pT3: intervention:13 (17) control:12 (15)</p> <p>pT4: intervention: 6 (8) control:4 (5)</p> <p>Lymph node stage, n (%)</p> <p>pN0: intervention:23 (29) control:25 (31)</p> <p>pN1: intervention:36 (46) control:39 (48)</p> <p>pN2: intervention:11 (14) control:9 (11)</p> <p>pN3: intervention:9 (11) control:8 (10)</p>	Manual lymph drainage (MLD)	No MLD	60	<p>1. Risk of lymphedema : defined as</p> <p>(1) ≥ 200 ml/ ≥ 2 cm increase of absolute arm volume difference compared with pre-surgical value by water displacement</p> <p>(2) ≥ 2 cm increase of arm circumference difference of olecranon at two adjacent measurement points compared with pre-surgical value</p> <p>(3) $\geq 5\%$/ $\geq 10\%$ increase of relative arm volume difference compared with pre-surgical value by water displacement</p> <p>(4) $\geq 5\%$ increase of relative arm volume difference compared with pre-surgical value by water displacement (persistent)</p> <p>2. Volume change : absolute change in difference in arm volume</p> <p>3. % volume change : relative change in difference in arm Volume</p>
--	--	-------------	-----------	-----	-----------	---------------	--	---	-----------------------------	--------	----	--

												<p>4. Change in physical activity : physical function (Lymph-ICF) and physical health (SF-36)</p> <p>5. Functional impairment-range of motion: shoulder flexion/abduction/external rotation/internal rotation</p>
Zhang, 2016 (Zhang et al., 2016)	RCT	1000(500/500)	NR	0	NR	Breast cancer	Surgery	Tumor-node-metastasis staging: I+II: control:211, intervention:197; III: control:289, intervention:303.	Self-manual lymph drainage (MLD) and exercise	Exercise	12	<p>1. Risk of lymphedema: by measuring upper limb circumferences on the surgical side compared with that of the contralateral side</p> <p>2. Functional impairment-range of motion: number of patients achieving 180 degrees of shoulder abduction in 6 months</p>

C. Physiotherapy with or without MLD vs. No physiotherapy

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Box, 2002 (Box, Reul-Hirche, Bullock-Saxton, & Furnival, 2002)	RCT	65 (32/33)	53.03/59	0	Median 24.3/27.2	Breast cancer	Surgical Procedure CLE&AD: control:51.5, intervention:46.9; MRM: control: 48.5, intervention:53.1. Radiotherapy control:48.5, interevrntion:65.6. Chemotherapy: control:21.2, interevention:43.8.	NR	Treatment group: received the Physiotherapy Management Care Plan (PMCP), including principles for lymphoedema risk minimization and early management of this condition when it was identified.	Control group: only received an exercise instruction booklet	24	Risk of lymphedema : defined by (1) an increase of ≥ 200 ml from the preoperative (VOL (OA-UOA) _{TA}). Arm volumes (VOL) were measured by water displacement. (2) an increase of ≥ 5 cm from the preoperative (CIRC (OA-UOA) _{TA}) (3) MFBIA ratio: number with ratio <95% lower reference limit (4) MFBIA % change: number with >10% decrease from the OA/UOA ratio

Box, 2002-2 (Box et al., 2002)	RCT (the same trial as Box, 2002)	65 (32/33)	53.03/59	0	Median 24.3/27.2	Breast cancer	Surgical Procedure CLE&AD: control:51.5, intervention:46.9; MRM: control: 48.5, intervention:53.1. Radiotherapy control:48.5, interevntion:65.6. Chemotherapy: control:21.2, interevention:43.8.	NR	Treatment group: Received the Physiotherapy Management Care Plan (PMCP), including principles for lymphoedema risk minimization and early management of this condition when it was identified.	Control group: only received an exercise instruction booklet	24	1. Change in physical activity : measured by compliance with the exercise protocol, which is associated with different patterns of recovery for all movements except external rotation. 2. Functional impairment-range of motion: shoulder abduction
-----------------------------------	-----------------------------------	------------	----------	---	---------------------	---------------	---	----	---	---	----	---

<p>Singh, 2013 (Singh et al., 2013)</p>	<p>Quasi Randomised</p>	<p>72 (41/31)</p>	<p>55.1/62.8</p>	<p>0</p>	<p>27.1/27.0</p>	<p>Breast cancer</p>	<p>All patients received surgery and some patients also received chemotherapy or radiation. Adjuvant radiation: intervention: 22 (53.7), control: 14 (45.2); Adjuvant chemotherapy: intervention: 16 (39), control:16 (32.3).</p>	<p>Cancer stage DCIS or Stage 1: intervention:2 (4.9), control:2 (6.5); Stage II: intervention:14 (34.1), control:10 (32.2); Stage III: intervention:19 (46.3), control:13 (41.9); Unknown: intervention: 6 (14.6), control:6 (19.4).</p>	<p>Clinical care pathway: preoperative education, prospective monitoring, and early physiotherapy</p>	<p>No clinical care pathway: preoperative education alone</p>	<p>7</p>	<p>1. Risk of lymphedema: a 2 cm increase in upper extremities circumference between adjacent points, compared with pre-surgical value 2. Change in physical activity: evaluated by Disabilities of Arm, Shoulder and Hand (DASH) 3. Functional impairment-range of motion: shoulder flexion/abduction/external rotation</p>
---	-------------------------	-------------------	------------------	----------	------------------	----------------------	---	---	---	---	----------	--

<p>Torres Lacomba, 2010 (Torres Lacomba et al., 2010)</p>	<p>RCT (ISRCTN9 5870846)</p>	<p>120 (60/60)</p>	<p>52.9/52.9</p>	<p>0</p>	<p>27.9/26.2</p>	<p>Breast cancer</p>	<p>Surgery- Quadrantectomy: Intervention: 24, Control: 26; Modified mastectomy: Intervention: 23, Control: 20; Lumpectomy: Intervention: 13, Control: 14. Postoperative therapy- Radiotherapy: intervention:44, control:49; Chemotherapy: interevntion: 50, control: 45; Hormonal therapy: Intervention: 39, Control: 33</p>	<p>NR</p>	<p>Physiotherapy program, including manual lymph drainage, massage of scar tissue, and progressive active and action assisted shoulder exercises. Also received the educational strategy.</p>	<p>Educational strategy</p>	<p>12</p>	<p>1. Risk of lymphedema: ≥2cm increase in the circumference of any two adjacent points compared with measurements in the other arm 2. % volume change: change in volume ratio</p>
---	--------------------------------------	--------------------	------------------	----------	------------------	----------------------	--	-----------	---	---------------------------------	-----------	--

D. Delayed vs. Early exercise

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Bendz, 2002 (Bendz & Fagevik Olsen, 2002)	RCT	205 (101/104)	58/58	0	NR	Breast cancer	All patients received surgery and some patients also received radiation. Mastectomy only: intervention:31, control:22; Incl. radiotherapy: intervention:5, control:7. Quadrant resection only: intervention:20, control:23; Incl. radiotherapy: intervention:45 control:52.	NR	Delayed shoulder exercise	Early shoulder exercise	24	1. Risk of lymphedema: volume on the operated side >10% increased compared with the non-operated side 2. % volume change: the arm volume measured volumetrically 3. Functional impairment-grip : grip strength measured with Vigortmeter 4. Functional impairment-range of motion : shoulder flexion/abduction/external rotation/internal rotation

Todd, 2008 (Todd et al., 2008)	RCT	116 (58/58)	56.5/57.2	0	27.5/28.4	Breast cancer	Surgery: wide local excision or mastectomy, including axillary lymph node dissection	Grade I intervention: 8, control: 8; Grade II intervention: 24, control: 27; Grade III intervention: 26, control: 23.	Intervention group (Delayed mobilisation): arm exercises and shoulder movement restricted to below shoulder level for the first 7 days after surgery.	Control group (Early mobilisation): commenced an exercise programme that incorporated exercises above shoulder level within 48 hours.	12	1. Risk of lymphedema: defined by a limb volume difference of 200 ml or more compared with the contralateral arm, measured using volume displacement 2. Functional impairment-grip : grip strength using a hand-held Jamar® dynamometer 3. Functional impairment-range of motion : shoulder abduction
-----------------------------------	-----	-------------	-----------	---	-----------	---------------	--	---	---	---	----	---

E. Preventive care/monitoring vs. Usual care

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Sato, 2014 (Sato et al., 2014)	Non-randomized controlled trial	149 (90/59)	ALND:52.9 SLNB:54.3/ ALND:52.1 SLNB:53.7	0	NR	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Radiotherapy: ALND intervention:69.2, control:83.3; SLNB intervention:58.8, control: 51.7. Chemotherapy, molecular targeting therapy: ALND intervention:79.5, control:86.7; SLNB intervention:11.8, control:24.1.	Stage 0: ALND intervention:0.0, control:6.7; SLNB intervention:25.5, control:44.8. Stage I: ALND intervention:7.7, control:20.0; SLNB intervention:54.9, control:48.3. Stage II: ALND intervention:43.6, control:50.0; SLNB intervention:17.6, control:6.9. Stage III: ALND intervention:41.0, control:23.3; SLNB intervention:2.0, control:0.0. Stage IV:ALND intervention:7.7, control:0.0; SLNB intervention:0.0, control:0.0.	Educational program: monitoring arm function and exercises	Routine care	3	1. Volume change: upper arm/forearm girth 2. Change in physical activity: evaluated by Disabilities of Arm, Shoulder and Hand (DASH) 3. Functional impairment-grip : grip strength 4. Functional impairment-range of motion : shoulder flexion/abduction/horizontal extension

Singh, 2013 (Singh et al., 2013)	Quasi Randomised	72 (41/31)	55.1/62.8	0	27.1/27.0	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Adjuvant radiation: intervention: 22 (53.7), control: 14 (45.2); Adjuvant chemotherapy: intervention: 16 (39), control:16 (32.3).	Cancer stage DCIS or Stage 1: intervention:2 (4.9), control:2 (6.5); Stage II: intervention:14 (34.1), control:10 (32.2); Stage III: intervention:19 (46.3), control:13 (41.9); Unknown: intervention: 6 (14.6), control:6 (19.4).	Clinical care pathway: preoperative education, prospective monitoring, and early physiotherapy	No clinical care pathway: preoperative education alone	7	1. Risk of lymphedema: a 2 cm increase in upper extremities circumference between adjacent points, compared with pre-surgical value 2. Change in physical activity: evaluated by Disabilities of Arm, Shoulder and Hand (DASH) 3. Functional impairment-range of motion: shoulder flexion/abduction/external rotation
Temur, 2019 (K. Temur & S. Kapucu, 2019)	RCT	61 (30/31)	47.6/45.6	0	>25: 19; <24.9: 11/ >25: 19; <24.9: 12	Breast cancer	Surgery: modified radical mastectomy and breast-conserving surgery	Stage 1 intervention: 2, control: 2; Stage 2 intervention: 16, control: 16; Stage 3 intervention: 12, control: 13.	Intervention group: Provided training and given a training booklet describing "exercise, massage and prevention methods"	Control group	6	1. Volume change : lateral epicondylitis (10 cm above/below), forearm (wrist joints 10 cm below) and metacarpophalangeal 2. Change in physical activity : physical functioning domain of EORTC QLQ-30

F. MLD vs. Exercise

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
de Oliveira, 2014 (de Oliveira et al., 2014)	Non-randomized controlled trial	96 (48/48)	55.6/56.7	0	28/29.1	Breast cancer	Surgery: Patey Modified Radical Mastectomy: MLD: 17, exercise: 27; Madden MRM: MLD: 25, exercise:19; Halsted RM: MLD: 1, exercise: 0; Neoadjuvant chemo: MLD: 29, exercise: 22.	Surgical staging: I: MLD: 0, exercise: 1; II: MLD: 9, exercise: 17; III/IV: MLD: 34, exercise: 28.	Manual lymphatic drainage (MLD)	Exercise	2	1. Volume change: hand, wrist, arm and forearm 2. Functional impairment-range of motion: shoulder flexion/abduction

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

G. Surveillance vs. No surveillance

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Yang, 2016 (Yang et al., 2016)	Comparative observational study	707 (390/317)	47.6/48.6	0	24.3/23.3	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Radiotherapy- Not done: intervention:47 (12.0 %), control:41 (12.9 %); Breast only: intervention:211 (54.0 %), control:174 (54.9 %); Breast and SCRT: intervention:140 (36.0 %), control:102 (32.2 %). Chemotherapy- Doxetaxel: intervention:280 (71.9 %), control:223 (70.3 %).	Histopathologic stage: I(%):intervention:109 (28.0 %), control:92 (29.0 %); II(%):intervention:215 (55.0 %), control:168 (53.0 %); III(%):intervention:66 (17.0 %), control:57 (18.0 %).	Surveillance program for lymphedema management (SLYM)	Standard care	60	Risk of lymphedema: defined by reference to the guidelines of the International Society of Lymphology (ISL) consensus document

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

H. Preoperative LS vs. No LS

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Boccardo, 2009 (Boccardo et al., 2009)	RCT	49 (25/24)	53.4/54.6	0	27.3/29.9	Breast cancer	Surgical Procedure (n) CLE & AD: intervention:12, control:13; MRM & AD: intervention:13, control:11. All included patients received radiotherapy.	NR	Preventive protocol (PG): Preoperative upper limb lymphscintigraphy (LS), principles for lymphedema risk minimization, and early management of this condition when it was identified.	Control group (CG)	24	1. Risk of lymphedema: a difference of over 200 ml from preoperative VOL measurements VOL (OA-UOA) [operated arm=OA; unoperated arm=UOA] 2. % volume change: arm volumes (VOL) were assessed using water displacement and measured to the nearest 5 ml

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Table S3. Certainty of evidence of the studies included in the meta-analysis

Programmed exercise with or without MLD vs. No intervention or usual care				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I-squared	Risk Difference	
Development of lymphedema	7 studies, 5 RCTs (791, events 159)	0.56 (0.35, 0.89), 59.8%	93 fewer per 1,000 (137 fewer to 23 fewer)	Low, rated down due to imprecision* and inconsistency [†]
	5 RCTs (among the 7 studies, 706, events 106)	0.77 (0.54, 1.10), 0%	40 fewer per 1,000 (79 fewer to 17 more)	Low, rated down twice due to imprecision*
Change in physical activity	2 studies, 2 RCTs (210)	3.54 (2.67, 4.41), 0%		Moderate, rated down due to imprecision*
Grip strength	1 study, 1 RCT (129)	-0.34 (-0.69, 0.01)		Low, rated down due to imprecision*
ROM – Shoulder flexion	3 studies, 3 RCTs (347)	9.36 (-2.36, 21.09), 92.3%		Very low, rated down twice due to imprecision* and inconsistency [†]
ROM – Shoulder abduction	3 studies, 3 RCTs (347)	11.38 (-6.65, 29.41), 94%		Very low, rated down twice due to imprecision* and inconsistency [†]
ROM – Shoulder external rotation	2 studies, 2 RCTs (290)	0.11 (-0.12, 0.34), 0%		Low, rated down twice due to imprecision*
ROM – Shoulder horizontal extension	2 studies, 2 RCTs (217)	3.12 (0.06, 6.18), 0%		Low, rated down due to imprecision* and risk of bias [#]
Compression vs. No compression				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I-squared	Risk Difference	
Development of lymphedema	1 study, 1 RCT (69)	0.80 (0.60, 1.07)	163 fewer per 1,000 (325)	Very low, rated down twice due to imprecision* and risk of bias [#]

			fewer to 57 more)	
MLD vs. No MLD				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I-squared	Risk Difference	
Development of lymphedema	3 studies, 3 RCTs (1,199, events 97)	0.49 (0.14, 1.71), 82.5%	49 fewer per 1,000 (83 fewer to 69 more)	Very low, rated down twice due to imprecision*, inconsistency+ and risk of bias#
Change in physical activity	2 studies, 2 RCTs (173)	-0.09 (-1.01, 0.83), 85%		Very low, rated down twice due to imprecision*, inconsistency+ and risk of bias#
ROM – Shoulder flexion	2 studies, 2 RCTs (199)	-22.71 (-69.31, 23.89), 99.5%		Very low, rated down twice due to imprecision*, inconsistency+ and risk of bias#
ROM – Shoulder abduction	2 studies, 2 RCTs (199)	-29.68 (-87.99, 28.64), 99.7%		Very low, rated down twice due to imprecision*, inconsistency+ and risk of bias#
ROM – Shoulder external rotation	1 study, 1 RCT (158)	0.48 (0.17, 0.80)		Low, rated down due to imprecision*
Physiotherapy with or without MLD vs. No physiotherapy				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I-squared	Risk Difference	
Development of lymphedema	3 studies, 2 RCTs (245)	0.31 (0.15, 0.64), 0%	152 fewer per 1,000 (187 fewer to 79 fewer)	Moderate, rated down due to imprecision*
ROM – Shoulder abduction	1 study, 0 RCT (72)	4.68 (-0.96, 10.32)		Very low, rated down twice due to imprecision*
ROM – Shoulder external rotation	1 study, 0 RCT (72)	0.33 (-0.14, 0.80)		Very low, rated down twice due to imprecision*
Delayed vs. Early exercise				
Outcomes	Studies, Design	Effect		Certainty in evidence

	(patients)	RR, WMD or SMD (95%CI), I-squared	Risk Difference	
Development of lymphedema	2 studies, 2 RCTs (321, events 31)	0.47 (0.23, 0.96), 0%	70 fewer per 1,000 (102 fewer to 5 fewer)	Low, rated down due to risk of bias [#]
Grip strength	2 studies, 2 RCTs (297)	-0.03 (-0.31, 0.26)		Very low, rated down due to imprecision* and risk of bias [#]
ROM – Shoulder flexion	1 study, 1 RCT (205)	-5.00 (-8.44, -1.56)		Low, rated down due to imprecision* and risk of bias [#]
ROM – Shoulder abduction	1 study, 1 RCT (205)	-6.00 (-13.41, 1.41)		Very low, rated down twice due to imprecision* and risk of bias [#]
ROM – Shoulder external rotation	1 study, 1 RCT (205)	0.09 (-0.19, 0.36)		Very low, rated down twice due to imprecision* and risk of bias [#]
Preventive care/monitoring vs. Usual care				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I-squared	Risk Difference	
Change in physical activity	3 studies, 1 RCTs (282)	1.00 (-0.32, 2.33), 96%		Very low, rated down twice due to imprecision* and inconsistency [†]
ROM – Shoulder flexion	2 studies, 0 RCT (221)	2.50 (-1.26, 6.26), 37.1%		Very low, rated down twice due to imprecision*
ROM – Shoulder abduction	2 studies, 0 RCT (221)	1.65 (-4.22, 7.53), 69.2%		Very low, rated down twice due to imprecision*
MLD vs. Exercise				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I-squared	Risk Difference	
ROM – Shoulder flexion	1 study, 0 RCT (89)	-2.00 (-9.07, 5.07)		Very low, rated down twice due to imprecision*

ROM – Shoulder abduction	1 study, 0 RCT (89)	-7.00 (-17.07, 3.07)		Very low, rated down twice due to imprecision*
Surveillance vs. No surveillance				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I- squared	Risk Difference	
Development of lymphedema	1 study, 0 RCT (203)	2.06 (1.54, 2.76)	413 more per 1,000 (210 more to 686 more)	Very low, rated down due to imprecision*
Preoperative LS vs. No LS				
Outcomes	Studies, Design (patients)	Effect		Certainty in evidence
		RR, WMD or SMD (95%CI), I- squared	Risk Difference	
Development of lymphedema	1 study, 1 RCT (49)	0.24 (0.06, 1.02)	253 fewer per 1,000 (313 fewer to 7 more)	Very low, rated down due to imprecision* and risk of bias#

*Imprecision: without known thresholds, we rated down once for either events < 300 or 95% CI including appreciable benefits and harm (binary outcomes), and patients number < 500 or 95% CI crossing 0 (continuous outcomes).

+Inconsistency: without known thresholds, we rated down for $I^2 > 50\%$ (binary outcomes) and $> 75\%$ (continuous outcomes).

#Risk of bias: For RCT, using Cochrane Collaboration Risk of Bias tool and evaluating according to the two most important domains that may cause bias: the “Random sequence generation” and “Blinding of outcome assessment”;
For non-RCT, using Newcastle-Ottawa quality assessment scale and evaluating according to the two most important domains that may cause bias: the “representative of patients, reported as a consecutive sample of patients” and “comparability of patients”.

Both RCT and non-RCT are evaluated as following: 2 domains of low risk – low risk

1 low risk plus 1 high risk or unclear/ two unclear – unclear

1 unclear plus 1 high risk / two high risk – high risk

When > 50% of studies are RCTs and no inconsistency, we started at high quality; otherwise started at low.

Table S4. Characteristics of included studies comparing Progressive resistance training (RRT) vs Usual care

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Ammitzbol 1, 2019 ¹	RCT (NCT02518477)	158 (82/76)	53/52	0	26.5/26.6	Breast cancer	Radiation+surgery (all patients) / some patients from each group also received chemotherapy Adjuvant: Control: 45 (59%) Intervention: 48 (59%) Neoadjuvant: Control: 21 (28%) Intervention: 25 (30%)	Histologic stage of malignancy, No. (%) 1: Control:16 (21) Intervention:12 (15) 2 :Control:35 (46) Intervention:48 (59) 3 :Control:18 (24) Intervention:15 (18)	Progressive resistance training (PRT)	Usual care	12	Change in physical activity: physical functioning with QLQ C-30 assessing QoL

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Ammitzbol 1, 2019 ²	RCT (NCT02518477)	158 (82/76)	53/52	0	26.5/26.6	Breast cancer	Radiation+surgery (all patients) / some patients from each group also received chemotherapy Adjuvant: Control: 45 (59%) Intervention: 48 (59%) Neoadjuvant: Control: 21 (28%) Intervention: 25 (30%)	Histologic stage of malignancy, No. (%) 1: Control:16 (21) Intervention:12 (15) 2 :Control:35 (46) Intervention:48 (59) 3 :Control:18 (24) Intervention:15 (18)	Progressive resistance training (PRT)	Usual care	12	1. Risk of lymphedema: >3% increased interlimb volume difference 2. % Volume change in interlimb volume difference 3. Functional impairment-grip : grip strength-ipsilateral/ contralateral side 4. Functional impairment-range of motion : ipsilateral/ contralateral shoulder flexion, ipsilateral/ contralateral shoulder abduction, ipsilateral/ contralateral shoulder external rotation

Table S5. Characteristics of included studies comparing Compression vs No compression

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Stuiver, 2013 ³	RCT	80 (41/39)	59/58	49/59	NR	Melanoma/Ur ogenital Cancer Intervention: 22/18, control: 23/17	Surgery with inguinal lymph node dissection	NR	Compression stockings	Patient education	12	Risk of lymphedema: lower limb/genital, defined as a 10% or greater increase from baseline in volume of the proximal or distal half of the thigh or the lower leg, using the standardized Kühnke's method of surface measurement

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Table S6. Characteristics of included studies comparing Manual lymph drainage (MLD) vs No MLD

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Cho, 2016 ⁴	RCT	41 (21/20)	46.6/50.7	0	NR	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Chemotherapy- Intervention:9 (42.9), control:11 (55). Radiotherapy- intervention:21 (100), control:19 (95).	Stage I: intervention:5 (23.8), control:12 (60) III: intervention:16 (76.2), control:8 (40).	Manual lymph drainage (MLD)+physical therapy (PT)	Physical therapy (PT)	1	1. Risk of lymphedema: ≥ 3 % volume increase from baseline in the affected upper limb 2. Volume change: arm volume calculated from circumference measurements taken at 4-cm intervals from the dorsum of the wrist to the axilla 3. Change in physical activity: EORTC QLQ-C30 and DASH 4. Functional impairment-range of motion: shoulder flexion/ abduction

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Devoogdt, 2011 ⁵	RCT	160 (79/81)	55.8/54.5	1/1	26.6/26.2	Breast cancer	<p>Surgery- Mastectomy: Intervention: 52, Control: 56; Breast conserving: Intervention: 27, Control: 25.</p> <p>Radiotherapy- IMC and medial supraclavicular: Intervention: 69, Control: 67.</p> <p>Radiotherapy- axilla: Intervention: 8, Control: 5.</p> <p>Chemotherapy: Intervention: 50, Control: 58.</p> <p>Neo-adjuvant chemotherapy: Intervention: 14, Control: 14.</p> <p>Trastuzumab: Intervention: 14, Control: 7.</p> <p>Endocrine treatment: Intervention: 55, Control: 66.</p>	<p>Tumour size: pT0: intervention: 1, control: 0; pT1: intervention: 21, control: 26; pT2: intervention: 38, control:39; pT3: intervention: 13, control: 12; pT4: intervention: 6, control: 4.</p>	Manual lymph drainage (MLD)	No MLD	12	<p>1. Risk of lymphedema : (1) ≥ 200 mL increase in arm volume up to the reference point 16 cm proximal to the olecranon by water displacement (2) ≥ 2 cm increase in arm circumference at the olecranon and 4, 8, 12, 16, and 20 cm above and under the olecranon (3) using author developed questionnaire</p> <p>2. Volume change : arm volume change</p> <p>3. Change in physical activity : SF-36 (QoL : physical health)</p>

Devoogdt, 2018 ⁶	RCT (the same trial as Devoogdt, 2011)	160 (79/81)	55.8/54.5	1/1	26.6/26.2	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Radiotherapy- intramammary chain, medial supraclavicular: intervention:69 (87), control:67 (83); axilla: intervention:8 (10), control:5 (6). Chemotherapy- all: intervention:50 (63), control:58 (72); neo-adjuvant: intervention:14 (18), control:14 (17). taxane-based: intervention:45 (57), control:46 (57).	Only for tumor size and node stage. Tumour size, n (%) pT0: intervention:1 (1) control:0 (0) pT1: intervention:21 (27) control:26 (32) pT2: intervention:38 (48) control:39 (48) pT3: intervention:13 (17) control:12 (15) pT4: intervention: 6 (8) control:4 (5) Lymph node stage, n (%) pN0: intervention:23 (29) control:25 (31) pN1: intervention:36 (46) control:39 (48) pN2: intervention:11 (14) control:9 (11) pN3: intervention:9 (11) control:8 (10)	Manual lymph drainage (MLD)	No MLD	60	1. Risk of lymphedema : defined as (1) ≥ 200 ml/ ≥ 2 cm increase of absolute arm volume difference compared with pre-surgical value by water displacement (2) ≥ 2 cm increase of arm circumference difference of olecranon at two adjacent measurement points compared with pre-surgical value (3) $\geq 5\%$ / $\geq 10\%$ increase of relative arm volume difference compared with pre-surgical value by water displacement (4) $\geq 5\%$ increase of relative arm volume difference compared with pre-surgical value by water displacement (persistent) 2. Volume change : absolute change in difference in arm volume 3. % volume change : relative change in difference in arm Volume
-----------------------------	--	-------------	-----------	-----	-----------	---------------	---	--	-----------------------------	--------	----	--

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
												<p>4. Change in physical activity : physical function (Lymph-ICF) and physical health (SF-36)</p> <p>5. Functional impairment-range of motion: shoulder flexion/abduction/external rotation/internal rotation</p>
Zhang, 2016 ⁷	RCT	1000(500/500)	NR	0	NR	Breast cancer	Surgery	Tumor-node-metastasis staging: I+II: control:211, intervention:197; III: control:289, intervention:303.	Self-manual lymph drainage (MLD) and exercise	Exercise	12	<p>1. Risk of lymphedema: by measuring upper limb circumferences on the surgical side compared with that of the contralateral side</p> <p>2. Functional impairment-range of motion: number of patients achieving 180 degrees of shoulder abduction in 6 months</p>

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Zimmerman, 2012 ⁸	RCT	67 (33/34)	60.3/58.6	0	25.6/24.3	Breast cancer	Breast conserving therapy-MLD: 20, control: 20; Modified mastectomy:-MLD: 13, control: 14; Endocrine therapy-MLD: 14, control: 19; Chemotherapy-MLD: 13, control: 15; Radiotherapy-MLD: 22, control: 25.	Stage I: MLD: 12 (36%), control: 11 (32%); Stage II: MLD: 15 (46%), control: 16 (47%); Stage III: MLD: 6 (18%), control: 7 (21%).	Manual lymphatic drainage (MLD)	No MLD, but with self-drainage	6	1. Risk of lymphedema: defined as >5% volume difference between the upper limbs 2. Volume change: ipsilateral upper limb

Table S7. Characteristics of included studies comparing Manual lymph drainage (MLD) vs Exercise

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
de Oliveira, 2014 ⁹	Non-randomized controlled trial	96 (48/48)	55.6/56.7	0	28/29.1	Breast cancer	Surgery: Patey Modified Radical Mastectomy: MLD: 17, exercise: 27; Madden MRM: MLD: 25, exercise:19; Halsted RM: MLD: 1, exercise: 0; Neoadjuvant chemo: MLD: 29, exercise: 22.	Surgical staging: I: MLD: 0, exercise: 1; II: MLD: 9, exercise: 17; III/IV: MLD: 34, exercise: 28.	Manual lymphatic drainage (MLD)	Exercise	2	1. Volume change: hand, wrist, arm and forearm 2. Functional impairment-range of motion: shoulder flexion/abduction

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Oliveira, 2018 ¹⁰	Comparative observational study	106 (53/53)	NR	0	NR	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Neoadjuvant chemotherapy: exercise:24 (45.3), MLD:36 (67.9); Adjuvant chemotherapy: exercise:8 (36.4), MLD:18 (62.1); Radiotherapy: exercise:16 (72.7), MLD:26 (86.7).	Clinical Stage I: exercise:1 (2.0) MLD:0 (0.0) II: exercise:17 (34.0) MLD:9 (18.0) III/IV: exercise:32 (64.0) MLD:43 (82.0) Surgical Stage I: exercise:2 (3.8) MLD:1 (1.9) II: exercise:20 (37.7) MLD:11 (20.8) III/IV: exercise:31 (58.5) MLD:41 (77.4)	Manual lymphatic drainage (MLD)	Active exercise (AE)	30	Functional impairment-range of motion: shoulder flexion/abduction

Table S8. Characteristics of included studies comparing Exercise vs No exercise

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Anderson, 2012 ¹¹	RCT	104 (52/52)	NR	0	NR	Breast cancer	Surgery- Lumpectomy only: Control: 25, Intervention: 23; Mastectomy: Control: 24, Intervention: 28. N/A Control: 3, Intervention: 1. Chemotherapy- Control: 31, Intervention: 31; Tamoxifen: Control: 23, Intervention: 26. Radiation therapy: Control: 36, Intervention: 31.	Stage I: intervention 25, control 26; II: intervention 19, control 21 III: intervention 8, control 4 N/A: intervention 0, control 1.	Exercise	Usual care	18	Volume change: mean arm volume change
Donmez, 2017 ¹²	RCT	52 (25/27)	48.6/49.5	0	NR	Breast cancer	Surgery (either MRM: Modified Radical Mastectomy or PS: Breast Protective Surgery)	NR	Physical activity program (PAP) and simple lymphatic drainage (SLD)	Usual care	1.5	1. Volume change: 5cm, 10cm, 15cm, 20cm over arm/forearm and mean metacarpophalangeal (MCP) circumference 2. Change in physical activity: DASH (self-rated questionnaire that measures upper extremity disability and symptoms)

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Cinar, 2008 ¹³	RCT	57 (27/30)	52.6/51.1	0	28.7/27.7	Breast cancer	Modified radical mastectomy 100%. Radiation therapy- Intervention: 10, Control: 14. Chemotherapy- Intervention: 23, Control: 29.	NR	Exercise: 15 sessions of individual rehabilitation program and continued with home-based physical activity program for 8 weeks	Control: only received a form including the exercises, which could be performed by themselves	6	<ol style="list-style-type: none"> 1. Risk of lymphedema 2. Volume change: midpoint of the third metacarpophalangeal joint and the ulnar styliod process (0 point) and at 10, 20, 30, and 40 cm proximally from the ulnar styliod process were marked, the circumferences at each marked point were summed to determine the total value and to calculate the difference between the operated and unoperated arm. 3. Functional impairment-range of motion : shoulder flexion/abduction/internal rotation/external rotation/adduction/extension

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Corrado, 2018 ¹⁴	Comparative observational study	30 (15/15)	55.8/54.7	0	NR	Breast cancer	Surgery: modified radical mastectomy and axillary lymph node dissection (ALND). Completed active breast cancer treatment with chemotherapy and radiation therapy.	NR	Home exercise program	No exercise	3	Risk of lymphedema : defined as ≥ 4 cm the difference between the sum of the 2 measurements among the affected and the unaffected arm, evaluated by measuring the arm circumference 10 cm above and 10 cm below the olecranon with a standard measuring tape

<p>Kilbreath, 2012¹⁵</p>	<p>RCT (Australian Clinical Trials Registry, ACTRN012606000050550)</p>	<p>160 (81/79)</p>	<p>53.5/51.6</p>	<p>0</p>	<p>26.2/26.5</p>	<p>Breast cancer</p>	<p>Surgery (100%) Mastectomy: Intervention: 48%, Control: 47%; Axillary node dissection: Intervention: 62%, Control: 58%. Chemotherapy: Taxane (Pacli or Docetaxel) Intervention: 0, Control: 5; Anthracycline Intervention: 27%, Control: 27%; Taxane and anthracycline Intervention: 40%, Control: 39%. Other Intervention: 1%, Control: 3%. Radiotherapy (%) Intervention: 79%, Control: 76%.</p>	<p>Intervention- Stage (I/II/III) (%) 17:44:38 Control- Stage (I/II/III) (%) 19:37:44</p>	<p>Exercise</p>	<p>No exercise</p>	<p>6</p>	<p>1. Risk of lymphedema : defined as (1) arm volume \geq10%, derived from circumferential measurements (2) >2 cm in 2 or more measures of interlimb circumference difference, measured at 10-cm intervals from the ulnar styloid to 40 cm proximally on both arms (3) exceeds Bioimpedance Spectroscopy ratio, determined by dividing the impedance from the nonaffected limb by that from the affected limb and this ratio was compared to previously established cut- offs: a ratio \geq1.139 when surgery was on the dominant side and a ratio \geq1.066 when surgery was on the non-dominant side was indicative of lymphedema 2. Functional impairment- range of motion: shoulder forward flexion/ abduction/ horizontal</p>
-------------------------------------	--	--------------------	------------------	----------	------------------	----------------------	--	---	-----------------	--------------------	----------	--

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
												extension/ external rotation
Kilbreath, 2013 ¹⁶	RCT (Australian Clinical Trials Registry, ACTRN012606000050550)	160 (81/79)	53.5/51.6	0	26.2/26.5	Breast cancer	Surgery (100%) Mastectomy: Intervention: 48%, Control: 47%; Axillary node dissection: Intervention: 62%, Control: 58%. Chemotherapy: Taxane (Pacli or Docetaxel) Intervention: 0, Control: 5; Anthracycline Intervention: 27%, Control: 27%; Taxane and anthracycline Intervention: 40%, Control: 39%. Other Intervention: 1%, Control: 3%. Radiotherapy (%) Intervention: 79%, Control: 76%.	Intervention- Stage (I/II/III) (%) 17:44:38 Control- Stage (I/II/III) (%) 19:37:44	Exercise	No exercise	15	Risk of lymphedema: ipsilateral upper limb, swelling was determined using bioimpedance spectroscopy with reference to previously established cut-offs for lymphedema

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Kilgour, 2008 ¹⁷	RCT	27 (16/11)	50.6/49.1	0	NR	Breast cancer	All patients received modified radical mastectomy and axillary node dissection.	NR	Home based exercise	Usual care	14 days	<ol style="list-style-type: none"> 1. Volume change: circumferential tape measurements of the forearm at 5 and 10 cm below the distal border of the lateral epicondyle of the humerus 2. Functional impairment-grip : grip strength using hand-grip dynamometry 3. Functional impairment-range of motion : shoulder flexion/abduction/external rotation using goniometric techniques

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Sagen, 2009 ¹⁸	RCT	204 (104/100)	54/55	0	NR	Breast cancer	Surgery- Breast ablation: Intervention: 46, Control: 51; Breast conserving: Intervention: 57, Control: 49. Radiotherapy- supraclavicular nodes: Intervention: 47, Control: 40. Radiotherapy-breast: Intervention: 78, Control: 73. Chemotherapy- Intervention: 42, Control: 38.	Histology grade 1: intervention: 25, control: 19; 2: intervention: 54, control: 56; 3: intervention: 24, control: 24.	Exercise: no activity restriction and supervised physical therapy program	No exercise: activity restriction and usual care	24	1. Risk of lymphedema: 10% increase in Voldiff between the affected arm and the control arm 2. Volume change: (1) difference between the volume of the affected arm and the volume of the control arm using the Simplified Water Displacement Instrument (SWDI) (2) change of affected arm volume
Schmitz, 2010 ¹⁹	RCT (NCT0019 4363)	154 (77/77)	54/56	0	27.5/28.6	Breast cancer	All patients had surgery with at least 2 lymph nodes removed. Chemotherapy: Intervention: 56, Control: 53; Radiation: Intervention: 59, Control: 58.	Cancer stage: Ductal carcinoma in situ- intervention: 1, control: 0; 1- intervention: 43, control:43 2- intervention: 8, control: 6; 3- intervention: 25, control: 28.	Weight lifting	No exercise	12	1. Risk of lymphedema: defined by $\geq 5\%$ increase in arm swelling, measured by water displacement 2. Change in physical activity: International Physical Activity Questionnaire (IPAQ), total physical activity (TPA) in MET-min/week

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Sisman, 2012 ²⁰	Comparative observational study	55 (45/10)	NR	Totally 2%	NR	Breast cancer	Surgery-Modified radical mastectomy: 53; Breast-conserving surgery:2. Radiotherapy 50 Gy in 25 days: 25	Stage I: 7 Stage II: 25 Stage III: 20 Stage IV: 3	Exercise	No exercise	6	Risk of lymphedema: measured 7.5 cm over and 7.5 cm below the elbow and the two arms were compared, defined as > 1.5 cm arm circumference measurements

Table S9. Characteristics of included studies comparing Delayed exercise vs Early exercise

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Bendz, 2002 ²¹	RCT	205 (101/104)	58/58	0	NR	Breast cancer	All patients received surgery and some patients also received radiation. Mastectomy only: intervention:31, control:22; Incl. radiotherapy: intervention:5, control:7. Quadrant resection only: intervention:20, control:23; Incl. radiotherapy: intervention:45 control:52.	NR	Delayed shoulder exercise	Early shoulder exercise	24	1. Risk of lymphedema: volume on the operated side >10% increased compared with the non-operated side 2. % volume change: the arm volume measured volumetrically 3. Functional impairment-grip : grip strength measured with Vigormeter 4. Functional impairment-range of motion : shoulder flexion/abduction/external rotation/internal rotation

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Todd, 2008 ²²	RCT	116 (58/58)	56.5/57.2	0	27.5/28.4	Breast cancer	Surgery: wide local excision or mastectomy, including axillary lymph node dissection	Grade I intervention: 8, control: 8; Grade II intervention: 24, control: 27; Grade III intervention: 26, control: 23.	Intervention group (Delayed mobilisation): arm exercises and shoulder movement restricted to below shoulder level for the first 7 days after surgery.	Control group (Early mobilisation): commenced an exercise programme that incorporated exercises above shoulder level within 48 hours.	12	1. Risk of lymphedema: defined by a limb volume difference of 200 ml or more compared with the contralateral arm, measured using volume displacement 2. Functional impairment- grip : grip strength using a hand-held Jamar® dynamometer 3. Functional impairment- range of motion : shoulder abduction

Table S10. Characteristics of included studies comparing Physiotherapy vs No physiotherapy

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Box, 2002 ²³	RCT	65 (32/33)	53.03/59	0	Median 24.3/27.2	Breast cancer	Surgical Procedure CLE&AD: control:51.5, intervention:46.9; MRM: control: 48.5, intervention:53.1. Radiotherapy control:48.5, interevrntion:65.6. Chemotherapy: control:21.2, interevention:43.8.	NR	Treatment group: received the Physiotherapy Management Care Plan (PMCP), including principles for lymphoedema risk minimisation and early management of this condition when it was identified.	Control group: only received an exercise instruction booklet	24	Risk of lymphedema : defined by (1) an increase of ≥ 200 ml from the preoperative (VOL (OA-UOA) _{TA}). Arm volumes (VOL) were measured by water displacement. (2) an increase of ≥ 5 cm from the preoperative (CIRC (OA-UOA) _{TA}) (3) MFBIA ratio: number with ratio $< 95\%$ lower reference limit (4) MFBIA % change: number with $> 10\%$ decrease from the OA/UOA ratio

Downloaded on 07-04-2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email pubpermissions@ons.org. ONS reserves all rights.

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Box, 2002 ²⁴	RCT (the same trial as Box 2002)	65 (32/33)	53.03/59	0	Median 24.3/27.2	Breast cancer	Surgical Procedure CLE&AD: control:51.5, intervention:46.9; MRM: control: 48.5, intervention:53.1. Radiotherapy control:48.5, interevrntion:65.6. Chemotherapy: control:21.2, interevention:43.8.	NR	Treatment group: Received the Physiotherapy Management Care Plan (PMCP), including principles for lymphoedema risk minimisation and early management of this condition when it was identified.	Control group: only received an exercise instruction booklet	24	1. Change in physical activity : measured by compliance with the exercise protocol, which is associated with different patterns of recovery for all movements except external rotation. 2. Functional impairment-range of motion: shoulder abduction

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Torres Lacomba, 2010 ²⁵	RCT (ISRCTN9 5870846)	120 (60/60)	52.9/52.9	0	27.9/26.2	Breast cancer	<p>Surgery-</p> <p>Quadrantectomy: Intervention: 24, Control: 26;</p> <p>Modified mastectomy: Intervention: 23, Control: 20;</p> <p>Lumpectomy: Intervention: 13, Control: 14.</p> <p>Postoperative therapy-</p> <p>Radiotherapy: intervention:44, control:49;</p> <p>Chemotherapy: interevntion: 50, control: 45;</p> <p>Hormonal therapy: Intervention: 39, Control: 33</p>	NR	<p>Physiotherapy program: including manual lymph drainage, massage of scar tissue, and progressive active and action assisted shoulder exercises. Also received the educational strategy.</p>	Educational strategy	12	<p>1. Risk of lymphedema: ≥2cm increase in the circumference of any two adjacent points compared with measurements in the other arm</p> <p>2. % volume change: change in volume ratio</p>

Table S11. Characteristics of included studies comparing Surveillance, education, or clinical care vs No intervention or standard care

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Boccardo, 2009 ²⁶	RCT	49 (25/24)	53.4/54.6	0	27.3/29.9	Breast cancer	Surgical Procedure (n) CLE & AD: intervention:12, control:13; MRM & AD: intervention:13, control:11. All included patients received radiotherapy.	NR	Preventive protocol (PG): Preoperative upper limb lymphscintigraphy (LS), principles for lymphedema risk minimization, and early management of this condition when it was identified.	Control group (CG)	24	1. Risk of lymphedema: a difference of over 200 ml from preoperative VOL measurements VOL (OA-UOA) [operated arm=OA; unoperated arm=UOA] 2. % volume change: arm volumes (VOL) were assessed using water displacement and measured to the nearest 5 ml

Downloaded on 07-04-2024. Single-user license only. Copyright 2024 by the Oncology Nursing Society. For permission to post online, reprint, adapt, or reuse, please email: pubpermissions@ons.org. ONS reserves all rights.

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Sato, 2014 ²⁷	Non-randomized controlled trial	149 (90/59)	ALND:52.9 SLNB:54.3/ ALND:52.1 SLNB:53.7	0	NR	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Radiotherapy: ALND intervention:69.2, control:83.3; SLNB intervention:58.8, control: 51.7. Chemotherapy, molecular targeting therapy: ALND intervention:79.5, control:86.7; SLNB intervention:11.8, control:24.1.	Stage 0: ALND intervention:0.0, control:6.7; SLNB intervention:25.5, control:44.8. Stage I: ALND intervention:7.7, control:20.0; SLNB intervention:54.9, control:48.3. Stage II: ALND intervention:43.6, control:50.0; SLNB intervention:17.6, control:6.9. Stage III: ALND intervention:41.0, control:23.3; SLNB intervention:2.0, control:0.0. Stage IV: ALND intervention:7.7, control:0.0; SLNB intervention:0.0, control:0.0.	Educational program: monitoring arm function and exercises	Routine care	3	1. Volume change: upper arm/forearm girth 2. Change in physical activity: evaluated by Disabilities of Arm, Shoulder and Hand (DASH) 3. Functional impairment-grip: grip strength 4. Functional impairment-range of motion: shoulder flexion/abduction/horizontal extension

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Singh, 2013 ²⁸	Quasi Randomised	72 (41/31)	55.1/62.8	0	27.1/27.0	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Adjuvant radiation: intervention: 22 (53.7), control: 14 (45.2); Adjuvant chemotherapy: intervention: 16 (39), control:16 (32.3).	Cancer stage DCIS or Stage 1: intervention:2 (4.9), control:2 (6.5); Stage II: intervention:14 (34.1), control:10 (32.2); Stage III: intervention:19 (46.3), control:13 (41.9); Unknown: intervention: 6 (14.6), control:6 (19.4).	Clinical care pathway: preoperative education, prospective monitoring, and early physiotherapy	No clinical care pathway: preoperative education alone	7	1. Risk of lymphedema: a 2 cm increase in upper extremities circumference between adjacent points, compared with pre-surgical value 2. Change in physical activity: evaluated by Disabilities of Arm, Shoulder and Hand (DASH) 3. Functional impairment-range of motion: shoulder flexion/abduction/external rotation
Temur, 2019 ²⁹	RCT	61 (30/31)	47.6/45.6	0	>25: 19; <24.9: 11/ >25: 19; <24.9: 12	Breast cancer	Surgery: modified radical mastectomy and breast-conserving surgery	Stage 1 intervention: 2, control: 2; Stage 2 intervention: 16, control: 16; Stage 3 intervention: 12, control: 13.	Intervention group: Provided training and given a training booklet describing "exercise, massage and prevention methods"	Control group	6	1. Volume change : lateral epicondylitis (10 cm above/below), forearm (wrist joints 10 cm below) and metacarpophalangeal 2. Change in physical activity : physical functioning domain of EORTC QLQ-30

Study Label (Author, Y)	Design*	Number of Patients (total, arm1/ arm 2)	Age in yr (mean, arm 1/ arm 2)	% Males	BMI (mean, arm 1/ arm 2)	Cancer type	Cancer treatment received	Cancer severity	First Arm (Intervention)	Second Arm (Control)	Follow up (Months)	Outcomes Reported
Yang, 2016 ³⁰	Comparative observational study	707 (390/317)	47.6/48.6	0	24.3/23.3	Breast cancer	All patients received surgery and some patients also received chemotherapy or radiation. Radiotherapy- Not done: intervention:47 (12.0 %), control:41 (12.9 %); Breast only: intervention:211 (54.0 %), control:174 (54.9 %); Breast and SCRT: intervention:140 (36.0 %), control:102 (32.2 %). Chemotherapy- Doxetaxel: intervention:280 (71.9 %), control:223 (70.3 %).	Histopathologic stage: I(%):intervention:109 (28.0 %), control:92 (29.0 %); II(%):intervention:215 (55.0 %), control:168 (53.0 %); III(%):intervention:66 (17.0 %), control:57 (18.0 %).	Surveillance program for lymphedema management (SLYM)	Standard care	60	Risk of lymphedema: defined by reference to the guidelines of the International Society of Lymphology (ISL) consensus document

References

1. Ammitzboll G, Kristina Kjaer T, Johansen C, et al. Effect of progressive resistance training on health-related quality of life in the first year after breast cancer surgery - results from a randomized controlled trial. *Acta Oncol.* 2019;1-8.
2. Ammitzboll G, Johansen C, Lanng C, et al. Progressive resistance training to prevent arm lymphedema in the first year after breast cancer surgery: Results of a randomized controlled trial. *Cancer.* 2019;11:11.
3. Stuiver MM, de Rooij JD, Lucas C, et al. No evidence of benefit from class-II compression stockings in the prevention of lower-limb lymphedema after inguinal lymph node dissection: results of a randomized controlled trial. *Lymphology.* 2013;46(3):120-131.
4. Cho Y, Do J, Jung S, Kwon O, Jeon JY. Effects of a physical therapy program combined with manual lymphatic drainage on shoulder function, quality of life, lymphedema incidence, and pain in breast cancer patients with axillary web syndrome following axillary dissection. *Support Care Cancer.* 2016;24(5):2047-2057.
5. Devoogdt N, Christiaens M-R, Geraerts I, et al. Effect of manual lymph drainage in addition to guidelines and exercise therapy on arm lymphoedema related to breast cancer: randomised controlled trial. *Bmj.* 2011;343:d5326.
6. Devoogdt N, Geraerts I, Van Kampen M, et al. Manual lymph drainage may not have a preventive effect on the development of breast cancer-related lymphoedema in the long term: a randomised trial. *J Physiother.* 2018;64(4):245-254.
7. Zhang L, Fan A, Yan J, et al. Combining Manual Lymph Drainage with Physical Exercise after Modified Radical Mastectomy Effectively Prevents Upper Limb Lymphedema. *Lymphat.* 2016;14(2):104-108.
8. Zimmermann A, Wozniewski M, Szklarska A, Lipowicz A, Szuba A. Efficacy of manual lymphatic drainage in preventing secondary lymphedema after breast cancer surgery. *Lymphology.* 2012;45(3):103-112.
9. de Oliveira MMF, de Rezende LF, do Amaral MTP, Pinto e Silva MP, Morais SS, Gurgel MSC. Manual lymphatic drainage versus exercise in the early postoperative period for breast cancer. *Physiother.* 2014;30(6):384-389.
10. Oliveira MMFd, Gurgel MSC, Amorim BJ, et al. Long term effects of manual lymphatic drainage and active exercises on physical morbidities, lymphoscintigraphy parameters and lymphedema formation in patients operated due to breast cancer: A clinical trial. *PLoS ONE.* 2018;13(1):e0189176.
11. Anderson RT, Kimmick GG, McCoy TP, et al. A randomized trial of exercise on well-being and function following breast cancer surgery: the RESTORE trial. *Journal of Cancer Survivorship.* 2012;6(2):172-181.
12. Donmez AA, Kapucu S. The effectiveness of a clinical and home-based physical activity program and simple lymphatic drainage in the prevention of breast cancer-related lymphedema: A prospective randomized controlled study. *Eur J Oncol Nurs.* 2017;31:12-21.
13. Cinar N, Seckin U, Keskin D, Bodur H, Bozkurt B, Cengiz O. The effectiveness of early rehabilitation in patients with modified radical mastectomy. *Cancer Nurs.* 2008;31(2):160-165.
14. Corrado B, Ciardi G, Iammarrone CS, Arpino G. Home Exercise Program is an effective tool in improving upper limb function and quality of life in breast cancer survivors: A retrospective observational study. *J Hum Sport Exerc.* 2018;13(4):926-939.
15. Kilbreath SL, Refshauge KM, Beith JM, et al. Upper limb progressive resistance training and stretching exercises following surgery for early

- breast cancer: a randomized controlled trial. *Breast Cancer Res Treat.* 2012;133(2):667-676.
16. Kilbreath SL, Lee M-J, Refshauge KM, et al. Transient swelling versus lymphoedema in the first year following surgery for breast cancer. *Support Care Cancer.* 2013;21(8):2207-2215.
 17. Kilgour RD, Jones DH, Keyserlingk JR. Effectiveness of a self-administered, home-based exercise rehabilitation program for women following a modified radical mastectomy and axillary node dissection: a preliminary study. *Breast Cancer Res Treat.* 2008;109(2):285-295.
 18. Sagen A, Karesen R, Risberg MA. Physical activity for the affected limb and arm lymphedema after breast cancer surgery. A prospective, randomized controlled trial with two years follow-up. *Acta Oncol.* 2009;48(8):1102-1110.
 19. Schmitz KH, Ahmed RL, Troxel AB, et al. Weight lifting for women at risk for breast cancer-related lymphedema: a randomized trial. *Jama.* 2010;304(24):2699-2705.
 20. Sisman H, Sahin B, Duman BB, Tanriverdi G. Nurse-assisted education and exercise decrease the prevalence and morbidity of lymphedema following breast cancer surgery. *J.* 2012;17(3):565-569.
 21. Bendz I, Fagevik Olsen M. Evaluation of immediate versus delayed shoulder exercises after breast cancer surgery including lymph node dissection--a randomised controlled trial. *Breast.* 2002;11(3):241-248.
 22. Todd J, Scally A, Dodwell D, Horgan K, Topping A. A randomised controlled trial of two programmes of shoulder exercise following axillary node dissection for invasive breast cancer. *Physiotherapy.* 2008;94(4):265-273.
 23. Box RC, Reul-Hirche HM, Bullock-Saxton JE, Furnival CM. Physiotherapy after breast cancer surgery: Results of a randomised controlled study to minimise lymphoedema. *Breast Cancer Research and Treatment.* 2002;75(1):51-64.
 24. Box RC, Reul-Hirche HM, Bullock-Saxton JE, Furnival CM. Shoulder movement after breast cancer surgery: results of a randomised controlled study of postoperative physiotherapy. *Breast Cancer Res Treat.* 2002;75(1):35-50.
 25. Torres Lacomba M, Yuste Sanchez MJ, Zapico Goni A, et al. Effectiveness of early physiotherapy to prevent lymphoedema after surgery for breast cancer: randomised, single blinded, clinical trial. *Bmj.* 2010;340:b5396.
 26. Boccardo FM, Ansaldi F, Bellini C, et al. Prospective evaluation of a prevention protocol for lymphedema following surgery for breast cancer. *Lymphology.* 2009;42(1):1-9.
 27. Sato F, Ishida T, Ohuchi N. The perioperative educational program for improving upper arm dysfunction in patients with breast cancer: a controlled trial. *Tohoku J Exp Med.* 2014;232(2):115-122.
 28. Singh C, De Vera M, Campbell KL. The effect of prospective monitoring and early physiotherapy intervention on arm morbidity following surgery for breast cancer: a pilot study. *Physiother Can.* 2013;65(2):183-191.
 29. Temur K, Kapucu S. The effectiveness of lymphedema self-management in the prevention of breast cancer-related lymphedema and quality of life: A randomized controlled trial. *Eur J Oncol Nurs.* 2019;40:22-35.
 30. Yang EJ, Ahn S, Kim E-K, et al. Use of a prospective surveillance model to prevent breast cancer treatment-related lymphedema: a single-center experience. *Breast Cancer Res Treat.* 2016;160(2):269-276.