

Translating Research on the Distress Thermometer Into Practice

Susan S. Tavernier, PhD, APRN-CNS, AOCN®



© Wavebreakmedia Ltd/Thinkstock

Distress is commonly experienced by people with cancer. The National Comprehensive Cancer Network (NCCN) developed a guideline for screening and managing distress in 1999 and has updated the guideline on an annual basis ever since. Using the five-phase framework for translational research, this article summarizes research findings related to the guideline and associated distress thermometer for each phase of translational research. The NCCN Distress Management Guideline is a useful tool to screen for and manage distress. However, additional research is needed, particularly in the areas of impact on outcomes of importance to patients.

Susan S. Tavernier, PhD, APRN-CNS, AOCN®, is an assistant professor in the School of Nursing at Idaho State University in Meridian. The author takes full responsibility for the content of the article. The author did not receive honoraria for this work. The content of this article has been reviewed by independent peer reviewers to ensure that it is balanced, objective, and free from commercial bias. No financial relationships relevant to the content of this article have been disclosed by the author, planners, independent peer reviewers, or editorial staff. Tavernier can be reached at tavesusa@isu.edu, with copy to editor at CJONEditor@ons.org. (Submitted September 2013. Accepted for publication September 15, 2013.)

Key words: distress; translational research; evidence-based practice

Digital Object Identifier: 10.1188/14.CJON.S1.26-30

Distress is a universal experience; however, the degree to which it is experienced, the duration, and the subsequent response to the distress varies widely. People living with cancer often have clinically relevant levels of distress requiring interventions for effective management (Carlson, Waller, Groff, Giese-Davis, & Bultz, 2013; Cohen, 2013; Kendall, Glaze, Oakland, Hansen, & Parry, 2011; Mitchell, Hussain, Grainger, & Symonds, 2011; Roerink et al., 2013; van't Spijker, Trijsburg, & Duivenvoorden, 1997; Zabora, BrintzenhofeSzoc, Curbow, Hooker, & Piantadosi, 2001; Zwahlen, Hagenbuch, Jenewein, Carley, & Buchi, 2011). Despite the widespread incidence of cancer-related distress, data indicate that clinicians are not familiar with valid and reliable tools to screen and assess for distress in their patients. In addition, the implementation of such tools in the clinical setting is limited (Jacobsen & Ransom, 2007; Pirl et al., 2007; Tavernier, Beck, & Dudley, 2013).

Assessment of cancer-related distress is best conducted using validated measures (Oncology Nursing Society, 2013). However, using an instrument congruent with the conceptual definition of distress is important. In addition, short, inexpensive measures easily completed by the patient, which also are incorporated into the electronic medical record, are ideal (Mitchell, Vahabzadeh, & Magruder, 2011). Although many instruments are used in research to measure distress, they are not always amenable to implementation in routine oncology practice. In addition, many instruments assess constructs related to distress

such as depression, symptom severity, or unmet needs, but not the global construct of distress. Several excellent reviews of the psychometric properties of such instruments are available (Mitchell, 2010; Vodermaier, Linden, & Siu, 2009; Ziegler et al., 2011), particularly the comprehensive data supplement to the review by Carlson, Waller, and Mitchell (2012).

The purpose of this article is to review research using the National Comprehensive Cancer Network® ([NCCN®], 2013) Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Distress Management and the distress thermometer. The American College of Surgeons (2012) Commission on Cancer will require routine screening of distress for all patients for healthcare agencies seeking accreditation beginning in 2015. Acknowledging the need for interdisciplinary collaboration for optimal management of distress, oncology nurses have the opportunity to demonstrate leadership in the selection, implementation, and evaluation of tools for assessing distress (Vitek, Rosenzweig, & Stollings, 2007). In addition, oncology nurses are well positioned to develop and evaluate interventions for improving outcomes related to distress management (Fitch, 2011).

The Distress Thermometer

The NCCN Guidelines for Distress Management was first published in 1999. The authors purposefully used the term *distress* to reduce the social stigma associated with other more

common terms associated with psychosocial problems (i.e., depression, anxiety, and maladjustment). The NCCN (2013) defines distress as

A multifactorial unpleasant emotional experience of a psychological (cognitive, behavioral, emotional), social, and/or spiritual nature that may interfere with the ability to cope effectively with cancer, its physical symptoms, and its treatment. Distress extends along a continuum, ranging from common normal feelings of vulnerability, sadness, and fears to problems that can become disabling, such as depression, anxiety, panic, social isolation, and existential and spiritual crisis (p. DIS-2).

The NCCN (2013) recommends screening for distress using a tool such as the distress thermometer, a single-item tool asking people to score how distressed they have been over the past week on a picture of a thermometer using a scale from 0 (no distress) to 10 (extreme distress) located at the top (see Figure 1). The person also completes a 39-item problem list, marking “yes” or “no” if the problem was experienced in the past week.

Overall, studies support the validity of the distress thermometer as a measure of cancer-related distress in adult and pediatric populations. Studies most often test construct validity of the distress thermometer with the Hospital Anxiety and Depression Scale (HADS), using HADS to determine specificity and sensitivity for identifying clinical cases. The distress thermometer was adapted and tested using English and Spanish versions in a pediatric oncology sample (Patel et al., 2011). A Danish version demonstrated validity in a study involving women with newly diagnosed breast cancer (Bidstrup et al., 2012). Validity is further supported in patients with advanced cancer (Ryan, Gallagher, Wright, & Cassidy, 2012) and family members of patients with cancer (Zwahlen et al., 2011). One limitation to validity studies is the predominance of studies occurring outside of the United States. Although a systematic review of cultural equivalence supports reliability, sensitivity, and specificity of the distress thermometer across languages and countries of study origin, Kayser, Acquati, and Tran (2012) identified the need for determining conceptual validity among immigrants and minority racial and ethnic groups within the United States.

Translation of the Distress Guideline Into Practice

Khuory et al. (2007) described five phases of translational research. The first phase (T0) focuses on a basic research question: “Do people with cancer experience distress?” T1 research is targeted toward a specific group: “What is the association between distress and people receiving cancer treatment?” The third translational research phase (T2) synthesizes research primarily for the goal of developing an evidence-based guideline. This level of research requires the use of valid measures of distress as well as support for interventions to decrease distress. The scope of the psychometric evidence is well presented in meta-analyses and systematic reviews of the measures used to assess distress and related concepts (i.e., anxiety and depression) (Carlson et al., 2012; Mitchell, 2010; Vodermaier et al., 2009; Ziegler et al., 2011).

Interventions to reduce distress are tested during the T2 phase of translational research. The NCCN Guidelines for Distress Management guideline provides a decision tree for referral based on the source of distress: physical, mental, social, spiritual, family, or practical problems. Therefore, interventions are targeted at the source of distress. Interventions to reduce distress related to physical symptoms are extensive and beyond the scope of this article, but more information can be found via the Oncology Nursing Society’s Putting Evidence Into Practice (PEP) guidelines, the NCCN’s Guidelines for Supportive Care, the Cochrane Review database, the National Guideline Clearinghouse, and the Joanna Briggs Institute, to name a few.

Interventions evaluating the efficacy of social interventions to reduce distress are inconclusive because of the poor quality of studies (Faller et al., 2013). Interventional research to alleviate cancer-related distress from spiritual sources is understudied. Published reports are primarily of qualitative design and in the area of palliative or end-of-life spiritual care. Care focusing on relationships and communication is viewed by patient participants as facilitators of spiritual care (Edwards, Pang, Shiu, & Chan, 2010).

A meta-analysis of randomized, controlled trials of psychological interventions for emotional distress concluded that interventions had a positive effect on reducing distress (Faller et al., 2013). The effect of the intervention was stronger in those with breast cancer or those who had metastatic disease. Most significantly, the investigators found that those interventions occurring during a longer period of time had a higher degree of sustained outcomes (Faller et al., 2013). Other studies support those conclusions (Carlson, Groff, Maciejewski, & Bultz, 2010; Franchi et al., 2013).

The fourth phase of translational research (T3) sometimes is referred to as dissemination, diffusion, or implementation research. In this stage, research focuses on using a guideline in the real-world setting. The research addresses the challenges of implementation because of the differences between the highly controlled environment in which research is conducted and the complexities of the clinical setting (Cochrane et al., 2007).

Three studies reported the diffusion of the NCCN Guidelines for Distress Management into practice. One study reported the uptake of the Guideline by NCCN institutions, with about half of the responding institutions routinely screening for distress, most using a tool other than the distress thermometer (Jacobsen & Ransom, 2007). A second survey of 448 oncologists found that 65% routinely assessed for distress, but only 14% used a validated instrument, and only 33% were at least somewhat familiar with the guideline (Pirl et al., 2007). The third study surveyed Oncology Nursing Society members (N = 420) working in ambulatory settings and reported that 38% of nurses were not at all familiar with the NCCN Guideline for Distress Management, and only 30% used a tool to assess for distress (Tavernier et al., 2013).

Suggestions have been made that distress should be considered as the “sixth vital sign” (Bultz & Carlson, 2005; Holland & Bultz, 2007). Feasibility of patients completing the distress thermometer digitally, using a tablet, is supported (Carlson et al., 2010), as is completion via telephone (Hughes, Sargeant, & Hawkes, 2011). The acceptability and usability of the distress thermometer adapted for pediatric patients experiencing

cancer was not supported because the source of distress often was unable to be determined, and nonsignificant differences in the quantity of services were noted (Patel et al., 2011).

Translational research studies examining the process or outcomes of implementing the NCCN Guideline for Distress Management or distress thermometer include evaluating the impact of screening on referrals, trajectories of distress, and barriers to adoption. An observational study cautiously concluded that when nurses were educated on the use of the distress thermometer, referral rates to psychosocial oncology services were higher and more accurate (Grassi et al., 2011). In a large (N = 505) Canadian study involving patients with a distress thermometer score greater than 4, investigators found that 24% of the patients accessed at least one support service; and, of the 24%, the majority accessed an average of three services (Waller, Williams, Groff, Bultz, & Carlson, 2013). Participants who were older, female, or had a lower education level were less likely to self-refer themselves despite high levels of distress. A study in Japan found that patients experiencing severe distress had sustained distress if they had high-intensity scores on pain, fatigue, or appetite loss when compared with those who had low distress (Yamaguchi et al., 2012). However, distress levels

were observed to be variable on repeated assessments, causing concern for over-referrals based on a single score (Carlson, Waller, Groff, Giese-Davis, et al., 2013; Yamaguchi et al., 2012).

Perceived barriers to adopting the distress thermometer or guideline include time, concern about inability to respond to increased demand for referrals, lack of knowledge about how to screen and manage distress, and no perceived benefit of screening (Absolom et al., 2011; Clark et al., 2012; Hughes et al., 2011; Oktay, Nedjat-Haiem, Davis, & Kern, 2012; Tavernier et al., 2013). Conversely, successful implementation, although reported primarily in case study or quality improvement projects, occurred when the implementation process had clear aims, regular data reporting, clinician comfort with introducing the distress thermometer, was perceived as beneficial to patient care, and was representative of team stakeholders in the planning and evaluation of the process (Absolom et al., 2011; Dudgeon et al., 2012; Hammonds, 2012; Oktay et al., 2012).

The fifth phase of translational research (T4) concerns the impact of a practice on populations. Because the goal of cancer treatment is to improve quality of life and/or increase longevity, an example of a research question related to the new American College of Surgeons standard is, “Does screening for distress

Instructions: First, please circle the number (0–10) that best describes how much distress you have been experiencing in the past week, including today.

Second, please indicate if any of the following has been a problem for you in the past week, including today. Be sure to check YES or NO for each.

YES	NO	Practical Problems	YES	NO	Physical Problems
<input type="checkbox"/>	<input type="checkbox"/>	Child care	<input type="checkbox"/>	<input type="checkbox"/>	Appearance
<input type="checkbox"/>	<input type="checkbox"/>	Housing	<input type="checkbox"/>	<input type="checkbox"/>	Bathing/dressing
<input type="checkbox"/>	<input type="checkbox"/>	Insurance/financial	<input type="checkbox"/>	<input type="checkbox"/>	Breathing
<input type="checkbox"/>	<input type="checkbox"/>	Transportation	<input type="checkbox"/>	<input type="checkbox"/>	Changes in urination
<input type="checkbox"/>	<input type="checkbox"/>	Work/school	<input type="checkbox"/>	<input type="checkbox"/>	Constipation
<input type="checkbox"/>	<input type="checkbox"/>	Treatment decisions	<input type="checkbox"/>	<input type="checkbox"/>	Diarrhea
		Family Problems	<input type="checkbox"/>	<input type="checkbox"/>	Eating
<input type="checkbox"/>	<input type="checkbox"/>	Dealing with children	<input type="checkbox"/>	<input type="checkbox"/>	Fatigue
<input type="checkbox"/>	<input type="checkbox"/>	Dealing with partner	<input type="checkbox"/>	<input type="checkbox"/>	Feeling swollen
<input type="checkbox"/>	<input type="checkbox"/>	Ability to have children	<input type="checkbox"/>	<input type="checkbox"/>	Fevers
<input type="checkbox"/>	<input type="checkbox"/>	Family health issues	<input type="checkbox"/>	<input type="checkbox"/>	Getting around
		Emotional Problems	<input type="checkbox"/>	<input type="checkbox"/>	Indigestion
<input type="checkbox"/>	<input type="checkbox"/>	Depression	<input type="checkbox"/>	<input type="checkbox"/>	Memory/concentration
<input type="checkbox"/>	<input type="checkbox"/>	Fears	<input type="checkbox"/>	<input type="checkbox"/>	Mouth sores
<input type="checkbox"/>	<input type="checkbox"/>	Nervousness	<input type="checkbox"/>	<input type="checkbox"/>	Nausea
<input type="checkbox"/>	<input type="checkbox"/>	Sadness	<input type="checkbox"/>	<input type="checkbox"/>	Nose dry/congested
<input type="checkbox"/>	<input type="checkbox"/>	Worry	<input type="checkbox"/>	<input type="checkbox"/>	Pain
<input type="checkbox"/>	<input type="checkbox"/>	Loss of interest in usual activities	<input type="checkbox"/>	<input type="checkbox"/>	Sexual
		Spiritual/Religious Concerns	<input type="checkbox"/>	<input type="checkbox"/>	Skin dry/itchy
			<input type="checkbox"/>	<input type="checkbox"/>	Sleep
			<input type="checkbox"/>	<input type="checkbox"/>	Substance abuse
			<input type="checkbox"/>	<input type="checkbox"/>	Tingling in hands/feet

Other Problems: _____

FIGURE 1. Distress Thermometer Screening Tool

Note. Adapted with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Distress Management V.2.2013. © 2013 National Comprehensive Cancer Network, Inc. All rights reserved. The NCCN Guidelines® and illustrations herein may not be reproduced in any form for any purpose without the express written permission of the NCCN. To view the most recent and complete version of the NCCN Guidelines, go online to NCCN.org. NATIONAL COMPREHENSIVE CANCER NETWORK®, NCCN®, NCCN GUIDELINES®, and all other NCCN Content are trademarks owned by the National Comprehensive Cancer Network, Inc.

Implications for Practice

- ▶ Use the distress thermometer from the National Comprehensive Cancer Network to assess patients for distress.
- ▶ Assess a patient's distress level on a routine basis because distress is influenced by many variables.
- ▶ Perform a follow-up assessment on patients reporting high levels of distress to ensure appropriate referral or intervention.

improve the quality of life and/or survival of patients with cancer?" The impact of screening with the distress thermometer on overall survival of cancer has not been explored. Screening using the distress thermometer followed by individual triage in patients with lung cancer led to fewer reports of pain and problems with coping and family conflict compared to those receiving usual care or screening alone (Carlson, Waller, Groff, & Bultz, 2013). Screening has been found to reduce the time to referral (Thewes, Butow, & Stuart-Harris, 2009).

Research is one component of evidence-based practice, with the primary goal being the improvement of patient care outcomes (Titler, 2011). Translational research related to the NCCN Guideline for Distress Management and distress thermometer exists for all five phases, albeit sparse for the T4 phase. The distress thermometer is a quick, valid indicator of the degree to which a patient is distressed by cancer and its treatment. Oncology nurses able to articulate the research evidence are better prepared to be engaged and assume a critical role in the decision making and implementation (Scott & McSherry, 2009) of an institution's approach to meet the Commission on Cancer standard.

Conclusion

Understanding that the distress thermometer is designed to screen for distress and is not a diagnostic tool is critical. Although the validity of the distress thermometer has been supported primarily using the HADS, the distress thermometer is not a diagnostic tool for depression or anxiety. When a patient is identified as having significant distress using the distress thermometer, additional assessment must occur to determine the source of distress. Additional assessment guides the appropriate referral or intervention. Using the distress thermometer should occur within those parameters and, as outlined in the NCCN Guidelines, a more in-depth assessment should be made for patients reporting levels of distress above established cutoff points.

Extensive research regarding the need to screen for cancer-related distress and the validity, feasibility, and acceptability of the distress thermometer exist. Although no significant intervention research findings for physical sources of distress have been noted, a need exists for research evaluating the efficacy of interventions related to the social, spiritual, and practical issues identified by the distress thermometer. Oncology nurses using the distress thermometer and the associated NCCN Guideline for Distress Management are encouraged to share the experience and to assist others attempting to integrate its use into routine care.

References

- Absolom, K., Holch, P., Pini, S., Hill, K., Liu, A., Sharpe, M., . . . Palliative Care Research. (2011). The detection and management of emotional distress in cancer patients: The views of health-care professionals. *Psycho-Oncology*, 20, 601–608.
- American College of Surgeons. (2012). *Cancer program standards 2012: Ensuring patient-centered care*. American College of Surgeons Commission on Cancer. Retrieved from <http://www.facs.org/cancer/coc/programstandards2012.pdf>
- Bidstrup, P.E., Mertz, B.G., Dalton, S.O., Deltour, I., Kroman, N., Kehlet, H., . . . Johansen, C. (2012). Accuracy of the Danish version of the "distress thermometer." *Psycho-Oncology*, 21, 436–443. doi:10.1002/pon.1917
- Bultz, B.D., & Carlson, L.E. (2005). Emotional distress: The sixth vital sign in cancer care. *Journal of Clinical Oncology*, 23, 6440–6441. doi:10.1200/jco.2005.02.3259
- Carlson, L.E., Groff, S.L., Maciejewski, O., & Bultz, B.D. (2010). Screening for distress in lung and breast cancer outpatients: A randomized controlled trial. *Journal of Clinical Oncology*, 28, 4884–4891. doi:10.1200/jco.2009.27.3698
- Carlson, L.E., Waller, A., Groff, S.L., & Bultz, B.D. (2013). Screening for distress, the sixth vital sign, in lung cancer patients: Effects on pain, fatigue, and common problems—Secondary outcomes of a randomized controlled trial. *Psycho-Oncology*, 22, 1880–1888. doi:10.1002/pon.3223
- Carlson, L.E., Waller, A., Groff, S.L., Giese-Davis, J., & Bultz, B.D. (2013). What goes up does not always come down: Patterns of distress, physical and psychosocial morbidity in people with cancer over a one-year period. *Psycho-Oncology*, 22, 168–176. doi:10.1002/pon.2068
- Carlson, L.E., Waller, A., & Mitchell, A.J. (2012). Screening for distress and unmet needs in patients with cancer: Review and recommendations. *Journal of Clinical Oncology*, 30, 1160–1177.
- Clark, P.G., Bolte, S., Buzaglo, J., Golant, M., Daratsos, L., & Loscalzo, M. (2012). From distress guidelines to developing models of psychosocial care: Current best practices. *Journal of Psychosocial Oncology*, 30, 694–714. doi:10.1080/07347332.2012.721488
- Cochrane, L.J., Olson, C.A., Murray, S., Dupuis, M., Tooman, T., & Hayes, S. (2007). Gaps between knowing and doing: Understanding and assessing the barriers to optimal health care. *Journal of Continuing Education in Healthcare Professions*, 27, 94–102. doi:10.1002/chp.106
- Cohen, M. (2013). The association of cancer patients' emotional suppression and their self-rating of psychological distress on short screening tools. *Behavioral Medicine*, 39, 29–35. doi:10.1080/08964289.2012.731440
- Dudgeon, D., King, S., Howell, D., Green, E., Gilbert, J., Hughes, E., . . . Sawka, C. (2012). Cancer Care Ontario's experience with implementation of routine physical and psychological symptom distress screening. *Psycho-Oncology*, 21, 357–364.
- Edwards, A., Pang, N., Shiu, V., & Chan, C. (2010). The understanding of spirituality and the potential role of spiritual care in end-of-life and palliative care: A meta-study of qualitative research. *Palliative Medicine*, 24, 753–770. doi:10.1177/0269216310375860
- Faller, H., Schuler, M., Richard, M., Heckl, U., Weis, J., & Küffner, R. (2013). Effects of psycho-oncologic interventions on emotional distress and quality of life in adult patients with cancer: Systematic review and meta-analysis. *Journal of Clinical Oncology*, 31, 782–793. doi:10.1200/jco.2011.40.8922

- Fitch, M.I. (2011). Screening for distress: A role for oncology nursing. *Current Opinion in Oncology*, 23, 331-337.
- Franchi, G., Bulli, F., Muraca, M.G., Maruelli, A., Grechi, E., & Miccinesi, G. (2013). Impact of a psycho-oncological rehabilitation intervention on psychological distress: The experience of CeRiOn (Oncological Rehabilitation Centre) Florence 2007-2010. *Supportive Care in Cancer*, 21, 2381-2386. doi:10.1007/s00520-013-1803-x
- Grassi, L., Rossi, E., Caruso, R., Nanni, M.G., Pedrazzi, S., Sofritti, S., & Sabato, S. (2011). Educational intervention in cancer outpatient clinics on routine screening for emotional distress: An observational study. *Psycho-Oncology*, 20, 669-674.
- Hammonds, L.S. (2012). Implementing a distress screening instrument in a university breast cancer clinic. *Clinical Journal of Oncology Nursing*, 16, 491-494.
- Holland, J.C., & Bultz, B.D. (2007). The NCCN guideline for distress management: A case for making distress the sixth vital sign. *Journal of the National Comprehensive Cancer Network*, 5, 3-7.
- Hughes, K.L., Sargeant, H., & Hawkes, A.L. (2011). Acceptability of the distress thermometer and problem list to community-based telephone cancer helpline operators, and to cancer patients and carers. *BMC Cancer*, 11, 46. doi:10.1186/1471-2407-11-46
- Jacobsen, P.B., & Ransom, S. (2007). Implementation of NCCN distress management guidelines by member institutions. *Journal of the National Comprehensive Cancer Network*, 5, 99-103.
- Kayser, K., Acquati, C., & Tran, T.V. (2012). No patients left behind: A systematic review of the cultural equivalence of distress screening instruments. *Journal of Psychosocial Oncology*, 30, 679-693. doi:10.1080/07347332.2012.721489
- Kendall, J., Glaze, K., Oakland, S., Hansen, J., & Parry, C. (2011). What do 1,281 distress screeners tell us about cancer patients in a community cancer center? *Psycho-Oncology*, 20, 594-600. doi:10.1002/pon.1907
- Khoury, M.J., Gwinn, M., Yoon, P.W., Dowling, N., Moore, C.A., & Bradley, L. (2007). The continuum of translation research in genomic medicine: How can we accelerate the appropriate integration of human genome discoveries into health care and disease prevention? *Genetic Medicine*, 9, 665-674. doi:10.1097/GIM.0b013e31815699d0
- Mitchell, A.J. (2010). Short screening tools for cancer-related distress: A review and diagnostic validity meta-analysis. *Journal of the National Comprehensive Cancer Network*, 8, 487-494.
- Mitchell, A.J., Hussain, N., Grainger, L., & Symonds, P. (2011). Identification of patient-reported distress by clinical nurse specialists in routine oncology practice: A multicentre UK study. *Psycho-Oncology*, 20, 1076-1083. doi:10.1002/pon.1815
- Mitchell, A.J., Vahabzadeh, A., & Magruder, K. (2011). Screening for distress and depression in cancer settings: 10 lessons from 40 years of primary-care research. *Psycho-Oncology*, 20, 572-584. doi:10.1002/pon.1943
- National Comprehensive Cancer Network. (2013). NCCN Clinical Practice Guidelines in Oncology for Distress Management [V.2.2013]. Retrieved from www.nccn.org/professionals/physician_gls/f_guidelines.asp#supportive
- Oktay, J.S., Nedjat-Haiem, F.R., Davis, C., & Kern, K.C. (2012). Distress screening: Experiences of oncology social workers. *Journal of Psychosocial Oncology*, 30, 652-666. doi:10.1080/07347332.2012.721490
- Oncology Nursing Society. (2013). Implementing screening for distress: The joint position statement from the American Psychosocial Oncology Society, Association of Oncology Social Work, and Oncology Nursing Society. *Oncology Nursing Forum*, 40, 423-424.
- Patel, S.K., Mullins, W., Turk, A., Dekel, N., Kinjo, C., & Sato, J.K. (2011). Distress screening, rater agreement, and services in pediatric oncology. *Psycho-Oncology*, 20, 1324-1333.
- Pirl, W.F., Muriel, A., Hwang, V., Kornblith, A., Greer, J., Donelan, K., . . . Schapira, L. (2007). Screening for psychosocial distress: A national survey of oncologists. *Journal of Supportive Oncology*, 5, 499-504.
- Roerink, S.H., de Ridder, M., Prins, J., Huijbers, A., de Wilt, H.J., Marres, H., . . . Netea-Maier, R.T. (2013). High level of distress in long-term survivors of thyroid carcinoma: Results of rapid screening using the distress thermometer. *Acta Oncologica*, 52, 128-137. doi:10.3109/0284186x.2012.723822
- Ryan, D.A., Gallagher, P., Wright, S., & Cassidy, E.M. (2012). Sensitivity and specificity of the distress thermometer and a two-item depression screen (Patient Health Questionnaire-2) with a "help" question for psychological distress and psychiatric morbidity in patients with advanced cancer. *Psycho-Oncology*, 21, 1275-1284.
- Scott, K., & McSherry, R. (2009). Evidence-based nursing: Clarifying the concepts for nurses in practice. *Journal of Clinical Nursing*, 18, 1085-1095. doi:10.1111/j.1365-2702.2008.02588.x
- Tavernier, S.S., Beck, S.L., & Dudley, W.N. (2013). Diffusion of a distress management guideline into practice. *Psycho-Oncology*. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1002/pon.3295/abstract;jsessionid=EC9BBB0467787B87E7D9A298A6B743E4.f03t01>
- Thewes, B., Butow, P., & Stuart-Harris, R. (2009). Does routine psychological screening of newly diagnosed rural cancer patients lead to better patient outcomes? Results of a pilot study. *Australian Journal of Rural Health*, 17, 298-304.
- Titler, M.G. (2011). Nursing science and evidence-based practice. *Western Journal of Nursing Research*, 33, 291-295.
- van't Spijker, A., Trijsburg, R.W., & Duivenvoorden, H.J. (1997). Psychological sequelae of cancer diagnosis: A meta-analytical review of 58 studies after 1980. *Psychosomatic Medicine*, 59, 280-293.
- Vitek, L., Rosenzweig, M.Q., & Stollings, S. (2007). Distress in patients with cancer: Definition, assessment, and suggested interventions. *Clinical Journal of Oncology Nursing*, 11, 413-418.
- Vodermaier, A., Linden, W., & Siu, C. (2009). Screening for emotional distress in cancer patients: A systematic review of assessment instruments. *Journal of the National Cancer Institute*, 101, 1464-1488. doi:10.1093/jnci/djp336
- Waller, A., Williams, A., Groff, S.L., Bultz, B.D., & Carlson, L.E. (2013). Screening for distress, the sixth vital sign: Examining self-referral in people with cancer over a one-year period. *Psycho-Oncology*, 22, 388-395. doi:10.1002/pon.2102
- Yamaguchi, T., Morita, T., Sakuma, Y., Kato, A., Kunimoto, Y., & Shima, Y. (2012). Longitudinal follow-up study using the distress and impact thermometer in an outpatient chemotherapy setting. *Journal of Pain and Symptom Management*, 43, 236-243.
- Zabora, J., BrintzenhofeSzoc, K., Curbow, B., Hooker, C., & Piantadosi, S. (2001). The prevalence of psychological distress by cancer site. *Psycho-Oncology*, 10, 19-28. doi:10.1002/1099-1611(200101/02)10:1<19::aid-pon501>3.0.co;2-6
- Ziegler, L., Hill, K., Neilly, L., Bennett, M.I., Higginson, I.J., Murray, S.A., & Stark, D. (2011). Identifying psychological distress at key stages of the cancer illness trajectory: A systematic review of validated self-report measures. *Journal of Pain and Symptom Management*, 41, 619-636.
- Zwahlen, D., Hagenbuch, N., Jenewein, J., Carley, M.I., & Buchi, S. (2011). Adopting a family approach to theory and practice: Measuring distress in cancer patient-partner dyads with the distress thermometer. *Psycho-Oncology*, 20, 394-403.